

# Recombinant Human CD99 protein, Fc-tagged

Cat. No. CD99-473H    Lot. No. (See product label)

## SPECIFICATION

<b>Product Overview</b>	Recombinant Human CD99 (Asp23-Asp122) protein was fused to human IgG1 Fc tag at C-terminus and expressed in human 293 cells (HEK293).
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Asp23-Asp122
<b>Description</b>	CD99 antigen is also known as MIC2, single-chain type-1 glycoprotein, HBA71, MIC2X, MIC2Y, MSK5X, 12E7, which belongs to the CD99 family. CD99/MIC2 is expressed on all leukocytes but highest on thymocytes. Involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. CD99/MIC2 plays a role in a late step of leukocyte extravasation helping leukocytes to overcome the endothelial basement membrane. CD99 acts at the same site as, but independently of PECAM1. There is also experimental evidence that it binds to cyclophilin A.
<b>Predicted N Terminal</b>	Asp23
<b>Form</b>	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.
<b>Molecular Mass</b>	The protein has a calculated MW of 36.2 kDa. The protein migrates as 45-50 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
<b>Endotoxin</b>	Less than 1.0 EU per µg by the LAL method.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

1/3

For Research Use Only

<b>Purity</b>	>90% as determined by SDS-PAGE.
<b>Storage</b>	<p>For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower.</p> <p>Please avoid repeated freeze-thaw cycles.</p> <p>This product is stable after storage at:</p> <ul style="list-style-type: none"> <li>-20 to -70 centigrade for 12 months in lyophilized state;</li> <li>-70 centigrade for 3 months under sterile conditions after reconstitution.</li> </ul>
<b>Reconstitution</b>	<p>It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.</p>

## GENE INFORMATION

<b>Gene Name</b>	CD99
<b>Official Symbol</b>	CD99
<b>Synonyms</b>	CD99; CD99 molecule; antigen identified by monoclonal antibodies 12E7, F21 and O13 , CD99 antigen , MIC2; CD99 antigen; E2 antigen; surface antigen MIC2; T-cell surface glycoprotein E2; MIC2 (monoclonal 12E7); antigen identified by monoclonal 12E7, Y homolog; antigen identified by monoclonal antibodies 12E7, F21 and O13; MIC2; HBA71; MIC2X; MIC2Y; MSK5X
<b>Gene ID</b>	4267
<b>mRNA Refseq</b>	NM_001122898
<b>Protein Refseq</b>	NP_001116370
<b>MIM</b>	313470


 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**UniProt ID**

P14209

**SDS-PAGE of CD99-473H**

Human CD99, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA

3/3

For Research Use Only