

Active Recombinant Human EPCAM, Fc-tagged, Biotinylated

Cat. No. EPCAM-571H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The recombinant human EPCAM-Fc fusion is expressed as a 470 amino acid protein consisting of Gln24 - Lys265 region of EPCAM (UniProt accession #P16422) and a C-terminal Fc from human IgG1, which exists as a dimer under non-reducing conditions.
Source	Human cells
Species	Human
Tag	Fc
Form	Supplied at 0.5 mg/ml in sterile PBS pH7.4 (carrier and preservative free). The purified recombinant protein was labeled with Biotin (3-5 Biotin per molecule).
Bio-activity	Binds anti-EPCAM monoclonal antibodies, human IgG1 and rabbit IgG.
Molecular Mass	Calculated molecular mass 53 kDa; estimated by SDS-PAGE under reducing condition 55-60 kDa probably due to glycosylation.
Endotoxin	<0.1 eu per 1 µg of purified recombinant protein determined by the
Purity	>95% judged by SDS-PAGE under reducing condition
Storage	The product is shipped at 4°C. Upon receipt, centrifuge the product briefly before opening the vial. It is recommended to store small aliquots at the temperature below -20°C for long-term storage and the product is stable for 3 months. The undiluted

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

protein can be stored at 4°C for no more than 2 weeks. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name EPCAM epithelial cell adhesion molecule [Homo sapiens]

Official Symbol EPCAM

Synonyms EPCAM; epithelial cell adhesion molecule; antigen identified by monoclonal AUA1 , M4S1, MIC18, TACSTD1, tumor associated calcium signal transducer 1; 17 1A; 323/A3; CD326; CO 17A; EGP 2; EGP34; EGP40; Ep CAM; ESA; GA733 2; HEA125; KS1/4; KSA; Ly74; MH99; MK 1; MOC31; TACST 1; TROP1; epithelial glycoprotein 314; human epithelial glycoprotein-2; cell surface glycoprotein Trop-1; adenocarcinoma-associated antigen; tumor-associated calcium signal transducer 1; major gastrointestinal tumor-associated protein GA733-2; membrane component, chromosome 4, surface marker (35kD glycoprotein); M4S1; MK-1; DIAR5; EGP-2; MIC18; EGP314; HNPCC8; TACSTD1; GA733-2;

Gene ID 4072


mRNA Refseq NM_002354

Protein Refseq NP_002345

MIM 185535

UniProt ID P16422

Chromosome Location 2p21



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

protein binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA