

Recombinant Human NGFR protein, His-tagged

Cat. No. NGFR-80H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human NGFR, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.
Description	NGFR, also known as tumor necrosis factor receptor superfamily member 16, is a member of the tumor necrosis factor receptor superfamily with a widespread pattern of expression in tissues such as the brain, liver, lung, and muscle. It plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin-dependent glucose uptake. They are low affinity receptor which can bind to NGF, BDNF, NT-3, and NT-4. It can mediate cell survival as well as cell death of neural cells.
Source	Insect cell
Species	Human
Tag	His
Form	Liquid. In Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.
Molecular Mass	24.6kDa (230aa)40-57kDa (SDS-PAGE under reducing conditions)
AA Sequence	KEACPTGLYT HSGECCKACN LGEGVAQPCG ANQTVCEPCL DSVTFSDVVS ATEPCKPCTE CVGLQSMSAP CVEADDAVCR CAYGYYQDETTGRCEACRVC EAGSGLVFSC QDKQNTVCEE CPDGTYSDEA NHVDPCLPCT VCEDTERQLR

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	ECTRWADAEC EEIPGRWITR STPPEGSDSTAPSTQEPEAP PEQDLIASTV AGVVTVMGS SQPVVTRGTT DNLEHHHHHH
Endotoxin	< 1.0 EU per 1 microgram of protein (determined by LAL method)
Purity	>95% by SDS – PAGE
Storage	Can be stored at +4 centigrade short term (1-2 weeks). For long term storage, aliquot and store at -20 centigrade or -70 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	1mg/ml (determined by Absorbance at 280nm)
GENE INFORMATION	
Gene Name	NGFR nerve growth factor receptor [<i>Homo sapiens</i>]
Official Symbol	NGFR
Synonyms	NGFR; nerve growth factor receptor; nerve growth factor receptor (TNFR superfamily, member 16); tumor necrosis factor receptor superfamily member 16; CD271; low affinity nerve growth factor receptor; p75NTR; TNFR superfamily; member 16; TNFRSF16; p75 ICD; NGF receptor; TNFR superfamily, member 16; low affinity neurotrophin receptor p75NTR; low-affinity nerve growth factor receptor; p75(NTR); Gp80-LNGFR;
Gene ID	4804
mRNA Refseq	NM_002507
Protein Refseq	NP_002498

MIM	162010
UniProt ID	P08138
Chromosome Location	17q21-q22
Pathway	Axonal growth inhibition (RHOA activation), organism-specific biosystem; Axonal growth stimulation, organism-specific biosystem; Cell death signalling via NRAGE, NRIF and NADE, organism-specific biosystem; Ceramide signalling, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; NADE modulates death signalling, organism-specific biosystem;
Function	death receptor activity; nerve growth factor binding; protein binding; receptor activity; signal transducer activity; transmembrane signaling receptor activity;