

Active Recombinant Human TGFB β 2, Fc-tagged, Biotinylated

Cat. No. TGFB β 2-697H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The recombinant human TGFB β 2-Fc fusion protein is expressed as a 365 amino acid protein consisting of Thr23 - Asp159 region of TGFB β 2 (UniProt accession #P37173) and a C-terminal Fc fusion from human IgG1, which exists as a dimer under non-reducing condition.
Source	Human cells
Species	Human
Tag	Fc
Form	Supplied at 0.5 mg/ml in sterile PBS pH7.4 (carrier & preservative free). The purified recombinant protein was labeled with Biotin (3-5 Biotin per molecule).
Bio-activity	Immobilized TGFB β 2 protein binds human TGF β 1 in a functional ELISA. Blocks TGF β 1-mediated signaling activity.
Molecular Mass	Calculated molecular mass (kDa): 41.1; Estimated by SDS-PAGE under reducing condition (kDa): ~60
AA Sequence	TIPPHVQKSVNNDMIVTDNNGAVKF PQLCKFCDVRFSTCDNQKSCMSNCS ITSICEKPQEVCAVWRKNDENIT LETVCHDPKLPYHDFILEDAAAPKC IMKEKKKPGETFFMCSSSDECNDN IIFSEEYNTSNPDSSGTHTCPPCP APELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVD

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GVEVHNAKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKALP
 APIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDI
 AVEWESNGQPENNYKTTTPVLDSGD SFFFLYSKLTVDKSRWQQGNVFCSCV
 MHEALHNHYTQKSLSLSPGK

Endotoxin <0.1 eu per 1 µg of purified recombinant protein determined by the lal>

Purity >95% judged by SDS-PAGE under reducing condition

Storage The product is shipped at 4°C. Upon receipt, centrifuge the product briefly before opening the vial. It is recommended to store small aliquots at the temperature below –20°C for long-term storage and the product is stable for 3 months. The undiluted protein can be stored at 4°C for no more than 2 weeks. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name **TGFB2** transforming growth factor, beta receptor II (70/80kDa) [Homo sapiens]

Official Symbol **TGFB2**

Synonyms TGFB2; transforming growth factor, beta receptor II (70/80kDa); MFS2, transforming growth factor, beta receptor II (70 80kD); TGF-beta receptor type-2; tbetaR-II; TGF-beta receptor type II; TGF-beta type II receptor; TGF-beta receptor type IIB; transforming growth factor-beta receptor type II; transforming growth factor beta receptor type IIC; transforming growth factor, beta receptor II (70/80kDa) isoform 1; transforming growth factor, beta receptor II (70/80kDa) isoform 2; AAT3; FAA3; MFS2; RIIC; LDS1B; LDS2B; TAAD2; TGFR-2; TGFbeta-RII;

Gene ID **7048**

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_001024847
Protein Refseq	NP_001020018
MIM	190182
UniProt ID	P37173
Chromosome Location	3p22
Pathway	ALK1 signaling events, organism-specific biosystem; Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chronic myeloid leukemia, organism-specific biosystem; Chronic myeloid leukemia, conserved biosystem;
Function	ATP binding; SMAD binding; glycosaminoglycan binding; metal ion binding; nucleotide binding; protein binding; contributes_to protein binding; receptor activity; receptor signaling protein serine/threonine kinase activity; transforming growth factor beta binding; transforming growth factor beta binding; transforming growth factor beta-activated receptor activity; transforming growth factor beta-activated receptor activity; transforming growth factor beta-activated receptor activity; transmembrane receptor protein serine/threonine kinase activity; type I transforming growth factor beta receptor binding; type I transforming growth factor beta receptor binding; type III transforming growth factor beta receptor binding;