

Recombinant Canine TWEAK (TNFSF12)

Cat. No. TWEAK-55C **Lot. No.** (See product label)

SPECIFICATION



Product Overview	Canine TWEAK/TNFSF12 was produced in yeast and therefore does not have endotoxin, is naturally folded, and post-translationally modified.
Description	Tumor necrosis factor ligand superfamily member 12 (TNFSF12) also known as TNF-related weak inducer of apoptosis (TWEAK) is a member of the tumor necrosis factor ligand (TNF) ligand family. Nineteen cytokines have been identified as part of the TNF family on the basis of sequence, functional, and structural similarities. Family members include TNF beta (TNFSF1), TNF alpha (TNFSF2), Lymphotoxin beta (TNFSF3), OX40 Ligand (TNFSF4), CD40 Ligand (TNFSF5), Fas Ligand (TNFSF6), CD27 Ligand (TNFSF7), CD30 Ligand (TNFSF8), 4-1BB Ligand (TNFSF9), TRAIL (TNFSF10), TRANCE/RANKL (TNFSF11), TWEAK (TNFSF12), APRIL(TNFSF13), BAFF (TNFSF13B), LIGHT (TNFSF14), TL1A/VEGI (TNFSF15), and GITR Ligand (TNFSF18). TWEAK has overlapping signaling functions with TNF, but displays a much wider tissue distribution. TWEAK can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. TWEAK is also found to promote proliferation and migration of endothelial cells, and thus acts as a regulator of angiogenesis.
Source	Yeast
Species	Canine
Form	Lyophilized
Molecular Mass	17.2 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	NASKGRKTRA RRAIAAHYEV HPQPGQDGGQ AGVDGTVSGW EEAKINSSNP LRYDRQSGEF IVTRAGLYYL YCQVHFDEGK AVYKLDLLV DDALALRCLE EFSATAASSL GPQLRLCQVS GLLPLRPGSS LRIRTLPAWH LKAAPFLTYF GLFQVN (156)
Applications	The Canine TWEAK/TNFSF12 protein can be used in cell culture, as a TWEAK ELISA Standard, and as a Western Blot Control.
Storage	-20 C
Tag	Non

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA