

HMGB1 ELISA Kit

Cat.No: DEIA6297V2

Lot. No. (See product label)

Size

96T

Intended use

CD provides a capture ELISA kit to determine HMGB1 levels in cell culture medium and sera. This kit contains enough reagents to measure 40 samples in duplicate together with standards.


General Description

HMGB1 (high mobility group box 1) (1) was recently rediscovered as a late lethal mediator of endotoxin (10) and is currently considered a pro-inflammatory cytokine that plays crucial roles in a variety of acute and chronic inflammatory diseases. HMGB1 contains 216 amino acids (6) that share more than 99% sequence identity in mice (2), rats (3), bovines (4), and humans (5). HMGB1 consists of three structural domains (7), termed "A box (9-85)", "B box (88-162)", and a negatively charged carboxyl terminus (186-216). Moreover, it has been previously shown that the B box recapitulates the pro-inflammatory activity whereas the A box acts as an antagonist of HMGB1 (8, 9).

Several lines of evidence indicate the significance of HMGB1 in the immune inflammatory response. For example, it has been shown that HMGB1 is actively released by a variety of cells such as macrophages when stimulated by lipopolysaccharides (LPS), TNF- α , and IL-1 (10), and is passively released by injured or necrotic cells associated with collapsing cell structures. In fact, patients who died from septic shock have higher serum HMGB1 levels than surviving sepsis patients (13). Similarly, high serum HMGB1 levels are observed in sepsis animal models and in collagen-induced arthritis animal models (14). With regard to the function of the protein itself, HMGB1 has also been shown to stimulate the release of TNF- α and IL-1 (11, 12), as well as bind LPS and synergistically increase peripheral blood mononuclear cell IL-6 production (19). Together, these observations demonstrate that HMGB1 plays important roles in the inflammatory cascade.

Assay Outline

HMGB1 Assay Outline



*Use one tip/sample. Do not cross-contaminate samples by re-using pipet tips. A multi-channel pipet is recommended.

Reagents And Materials Provided

1. HMGB1 Standard: 2 vials, 50 μ l/vial
2. Capture Antibody (Anti-HMGB1 Monoclonal Antibody): 1 vial, 100 μ l/vial
3. Detection Antibody (Anti-HMGB1 Monoclonal Antibody): 1 vial, Lyophilized
4. Solution A - Capture Antibody Dilution Buffer: 1 bottle, 10 ml
5. Solution B - Sample/Standard Dilution Buffer: 1 bottle, 20 ml
6. Solution C - Detection Antibody Dilution Buffer: 1 bottle, 10 ml
7. Solution D - Streptavidin Peroxidase Dilution Buffer: 1 bottle, 20 ml
8. Streptavidin Peroxidase: 2 vials, 50 μ l
9. TMB Solution (contains DMSO): 2 vials, 0.2 ml
10. Chromogen Dilution Buffer: 1 bottle, 20 ml
11. Stop Solution - 2N Sulfuric Acid: 1 bottle, 10 ml
12. Wash Buffer, 20X: 1 bottle, 50 ml
13. ELISA Plate: 1 each, 96-well (8-well strips x 12)


Storage

-20°C

Assay Procedure

All reagents must be at room temperature before use.

1. Add Capture Antibody: Dilute 1 vial (100 μ l) of Capture Antibody with 10 ml of Capture Antibody Dilution Buffer (Solution A). Add 100 μ l of capture antibody solution to each well and incubate at 4°C overnight. If you plan to use less, the remaining stock solution can be stored in its original vial at -20°C.
2. Prepare Standard Dilutions: The recommended standard range is 0.8-50 ng/ml. Dilute one vial of HMGB1 Standard with 950 μ l of Sample/Standard Dilution Buffer (Solution B) - 100 ng/ml. Prepare serial dilutions of the standard by mixing 200 μ l of the 100 ng/ml standard with 200 μ l of Solution B - 50 ng/ml. Then repeat this procedure to make six more serial dilutions of standard - 25, 12.5, 6.25, 3.1, 1.6, and 0.8 ng/ml solutions. Partially used 100 ng/ml standard stock cannot be saved for future assays. Discard unused, diluted standard solution. We recommend making fresh standard and serial dilutions for each assay.




3. Prepare Sample Dilutions: Centrifuge samples at 10,000 rpm at 4°C for 3 minutes to remove insoluble materials and lipids, and use the supernatant as samples. Dilute sample, at minimum, with an equal volume of Solution B. For example, take 100 µl of a serum, and mix with 100 µl of Solution B. If the HMGB1 level is higher than 50 ng/ml, re-assay the sample at a higher dilution.

4. Prepare Detection Antibody: Reconstitute one vial of Detection Antibody with 50 µl of distilled water to make a stock solution. Dissolve the 50 µl of detection antibody stock solution in 5 ml of Detection Antibody Dilution Buffer (Solution C). If you plan to use less, the remaining stock solution can be stored in its original vial at -20°C.

5. Dilute Wash Buffer: Dilute 50 ml of 20X wash buffer in 950 ml of distilled water (1X wash buffer). Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. Do not allow the plate to dry out.

6. Add Standards, Samples, and Detection Antibody: Mix standards, samples, and detection antibody tubes well. Add 50 µl of Solution B (blank), standards, and samples to appropriate wells (Figure 1). Add 50 µl of diluted detection antibody solution to all wells. Mix all wells by pipetting or use a plate shaker. Cover the plate with a plate sealer and incubate at 37°C for 1 hour, then transfer plate to 4°C overnight.

Figure 1 - A typical assay layout



7. Prepare Streptavidin Peroxidase: Dilute one vial (50 µl) Streptavidin Peroxidase in 10 ml of Streptavidin Peroxidase Dilution Buffer (Solution D). If you plan to use less, the remaining stock solution can be stored in its original vial at -20°C.

8. Wash: Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. Do not allow the plate to dry out.

9. Add Streptavidin Peroxidase: Add 100 µl of streptavidin peroxidase solution to each well and incubate at room temperature for 30 minutes.

10. Wash: Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. Do not allow the plate to dry out.

11. TMB: Use new tubes when preparing TMB. Dilute one vial (200 µl) TMB with 10 ml of Chromogen Dilution Buffer just prior to use. Add 100 µl of TMB solution to all wells immediately after washing the plate. Incubate for 30 minutes at room temperature. If you plan to use less, the remaining stock solution can be stored in its original vial at -20°C.


12. Stop: Add 50 µl of 2N sulfuric acid (Stop Solution) to each well.

13. Read Plate: Read the OD values at 450 nm (a 630 nm filter can be used as a reference) immediately. If the OD values of samples are greater than the OD values of the highest standard, re-assay the samples at a higher dilution.

Calculation

1. Average the duplicate OD values for the blank, standards, and samples.
2. Subtract the averaged blank (B) OD value from the averaged standard and sample OD values.
3. Plot the OD values of standards against the amount of HMGB1 (ng/ml) using a log scale. Figure 2 shows a typical standard curve where the HMGB1 range is from 0.8-50 ng/ml.
4. The concentration of HMGB1 (ng/ml) in samples can be calculated using regression analysis
- . Multiply the results by the dilution factors (usually 2 without extra dilution).

Figure 2 - A typical standard curve



Precision

Human Sera

Test At	2 ng/ml	7.5 ng/ml	30 ng/ml
Inter-Assay CV (%)	1.3	5.1	3.1
Intra-Assay CV (%)	2.7	3.4	3.6

Mouse Sera

Test At	2 ng/ml	7.5 ng/ml	30 ng/ml
Inter-Assay CV (%)	3.7	4.5	0.9
Intra-Assay CV (%)	1.4	3.2	8.1

Specificity

Average cross reactivity with bovine HMGB2 is 12.4%.

Recovery

Species	Averaged Recovery Results
Human (5 sera)	101 ± 25.1%
Mouse (5 sera)	104 ± 8.3%

Precautions

1. It is recommended that the standard and samples be run in duplicate.
2. Warm up all buffers to room temperature before use.
3. For partial reagent use, use the appropriate dilution ratio for each reagent as stated in the protocol. For example, if the protocol dilutes 50 μ l of stock solution in 10 ml of buffer, then for 6 strips, dilute 25 μ l of stock solution in 5 ml of buffer. Partially used capture, detection, streptavidin peroxidase, and TMB reagents may be kept at in their original vials at -20°C.
4. Crystals may form in the 20X wash buffer when stored at cold temperatures. If crystals have formed, it is necessary to warm the wash buffer by placing the bottle in warm water until crystals have dissolved completely.
5. Measure exact volumes of buffers using a serological pipette, as extra buffer is provided.
6. Cover the plate with plastic wrap or a plate sealer after each step to prevent evaporation from the outside wells of the plate.
7. This kit can be used to determine HMGB1 in serum and cell culture medium. However, special concern should be considered for assaying HMGB1 in human serum, because autoantibodies to HMGB1 are determined in 9-89% of sera from patients with autoimmune and inflammatory diseases (15-18). These reports indicate that human serum polyclonal antibodies to HMGB1 might mask the epitopes recognized by the capture and detection antibodies used in this kit, resulting in interference against the assay. Therefore, it is important to use this kit with background knowledge of patients.
8. This kit contains components of animal origin from non-infectious animals, but should be treated as potential biohazards in use and for disposal.
9. Human samples should be handled as if they are capable of transmitting infectious diseases.

References

1. Einck L and Bustin M. The intracellular distribution and function of the high mobility group chromosomal proteins. *Exp Cell Res.* 156:295- 310 (1985).
2. Ferrari S et al. The mouse gene coding for high mobility group 1 protein (HMG1). *J Biol Chem.* 269:28803-28808 (1994).
3. Paonessa G et al. Nucleotide sequence of rat liver HMG1 cDNA. *Nucleic Acids Res.* 15:9077 (1987).
4. Kaplan DJ and Duncan CH. Full length cDNA sequence for bovine high mobility group 1 (HMG1) protein. *Nucleic Acids Res.* 16 : 10375 (1988).
5. Wen L et al. A human placental cDNA clone that encodes nonhistone chromosomal protein HMG-1. *Nucleic Acids Res.* 17:1197-1214 (1989).
6. Parkkinen J et al. Amphoterin, the 30-kDa protein in a family of HMG1-type polypeptides. Enhanced expression in transformed cells, leading edge localization, and interactions with plasminogen activation. *J Biol Chem.* 268:19726-38 (1993).
7. Li J et al. Recombinant HMGB1 with cytokine-stimulating activity. *J Immunol Methods.* 289:211-23 (2004).
8. Bustin M. Regulation of DNA-dependent activities by the functional motifs of the high-mobility-group chromosomal proteins. *Mol Cell Biol.* 19:5237-46 (1999).
9. Yang H et al. Reversing established sepsis with antagonists of endogenous high-mobility group box 1. *Proc Natl Acad Sci U S A.* 101:296- 301 (2004).
10. Wang H et al. HMG-1 as a late mediator of endotoxin lethality in mice. *Science.* 285:248-251 (1999).
11. Wang H et al. Proinflammatory cytokines (tumor necrosis factor and interleukin-1) stimulate release of high mobility group protein-1 by pituicytes. *Surgery.* 126:389-392 (1999).
12. Abraham E et al. HMG-1 as a mediator of acute lung inflammation. *J Immunol.* 165:2950-2954 (2000).
13. Andersson U et al. High mobility group 1 protein (HMG-1) stimulates proinflammatory cytokine synthesis in human monocytes. *J Exp Med.* 192:565-570 (2000).
14. Kokkola R et al. Successful treatment of collagen-induced arthritis in mice and rats by targeting extracellular high mobility group box chromosomal protein 1 activity. *Arthritis Rheum.* 48:2052-2058 (2003).
15. Wittemann B et al. Autoantibodies to nonhistone chromosomal proteins HMG-1 and HMG-2 in sera of patients with juvenile rheumatoid arthritis. *Arthritis Rheum.* 33(9):1378-83 (1990).
16. Uesugi H et al. Prevalence and characterization of novel pANCA, antibodies to the high mobility group non-histone chromosomal proteins HMG1 and HMG2, in systemic rheumatic diseases. *J Rheumatol.* 25(4):703-9 (1998).
17. Sobajima J et al. High mobility group (HMG) non-histone chromosomal proteins HMG1 and HMG2 are significant target antigens of perinuclear anti-neutrophil cytoplasmic antibodies in autoimmune hepatitis. *Gut.* 44(6):867-73 (1999).
18. Sobajima J et al. Novel autoantigens of perinuclear anti-neutrophil cytoplasmic antibodies (P-ANCA) in ulcerative colitis: non-histone chromosomal proteins, HMG1 and HMG2. *Clin Exp Immunol.* 107(1):135-40 (1997).
19. Hreggvidsdottir HS et al. The alarmin HMGB1 acts in synergy with endogenous and exogenous danger signals to promote inflammation. *J Leukoc Biol.* 86(3):655-62 (2009).