



Anti-CD40 monoclonal antibody, clone LOB7/6 [R-PE] (CABT-46036MH)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Mouse anti Human CD40 antibody, clone LOB7/6 recognizes the human CD40 cell surface antigen, a 48kDa glycoprotein expressed by B lymphocytes and weakly by some monocytes. CD40 is involved in the process of B cell selection in germinal centres and is vital in T cell-B cell interactions. Flow Cytometry Use 10ul of the suggested working dilution to label 106 cells in 100ul.
Specificity	CD40
Immunogen	CD40 Ig(Fc) fusion protein containing the EC region of human CD40 and Fc region of human IgG
Isotype	IgG2a
Source/Host	Mouse
Species Reactivity	Human, Dog
Clone	LOB7/6
Conjugate	PE
Applications	FC
Format	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilised
Size	100 tests
Preservative	0.09% Sodium Azide
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected

from light. Should this product contain a precipitate we recommend microcentrifugation before use.

GENE INFORMATION

Gene Name	CD40 CD40 molecule, TNF receptor superfamily member 5 [Homo sapiens (human)]
Official Symbol	CD40
Synonyms	CD40; CD40 molecule, TNF receptor superfamily member 5; p50; Bp50; CDW40; TNFRSF5; tumor necrosis factor receptor superfamily member 5; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; CD40 antigen (TNF recept
Entrez Gene ID	958
Protein Refseq	NP_001241
UniProt ID	P25942
Chromosome Location	20q12-q13.2
Pathway	Adaptive Immune System; Allograft Rejection; Allograft rejection; Asthma; Autoimmune thyroid disease; CD40/CD40L signaling; Cell adhesion molecules (CAMs); Cytokine-cytokine receptor interaction;
Function	antigen binding; enzyme binding; protein binding; receptor activity; signal transducer activity; ubiquitin protein ligase binding;