



Anti-Cd339 monoclonal antibody, clone HMJ1-29 (CABT-49800HM)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview

Hamster anti Mouse CD339 antibody, clone HMJ1-29 specifically recognizes CD339, otherwise known as Jagged1, one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4. The Notch signaling pathway is an evolutionarily conserved pathway in multi-cellular organisms, which is vital for cell-cell communication, important during fundamental developmental and physiological processes, including regulation of cell fate decisions during neuronal, cardiac and endocrine development, stem cell hematopoiesis, thymic T-cell development, and both tumor progression and suppression. Ligation of Notch receptors by their specific ligands, Jagged1 (CD339), Jagged2, Delta-like protein 1 (DLL1), DLL3 and DLL4, on physically adjacent signal receiving cells, induces proteolysis of the receptors by ADAM-family metalloproteases and the gamma-secretase complex, within the transmembrane domain, releasing the Notch intracellular domain (NICD) to translocate to the nucleus. Subsequent signal transduction then occurs through either the CSL-NICD-Mastermind complex cascade (canonical pathway), or NF-kappaB-NICD and CSL-NICD-Deltex complex signaling cascades (non-canonical pathway). The canonical pathway inhibits the differentiation of stem cells or progenitor cells, whilst the non-canonical pathway promotes differentiation. Jagged1 signaling is implicated in cell-fate decisions during hematopoiesis, as well as in both early and late stages of mammalian cardiovascular development, and is involved in the inhibition of myoblast differentiation. Studies have shown a significant increase in the expression of Jagged1 in metastatic prostate cancer, compared with localized prostate cancer or benign prostatic tissues, implicating Jagged1 as a biomarker to facilitate their differentiation. In humans, mutations in the JAG1 gene are responsible for the autosomal dominant Flow Cytometry Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

Specificity	CD339
Immunogen	Jagged1-expressing CHO cells.
Isotype	IgG

Source/Host	Hamster
Species Reactivity	Mouse, Rat
Clone	HMJ1-29
Conjugate	Unconjugated
Applications	IHC-Fr; FC
Format	Purified IgG - liquid
Size	100 µg
Preservative	See individual product datasheet
Storage	in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

GENE INFORMATION

Gene Name	Jag1 jagged 1 [Mus musculus (house mouse)]
Official Symbol	JAG1
Synonyms	JAG1; jagged 1; Htu; Ozz; ABE2; Ser-1; Gsfabe2; protein jagged-1; Serrate-1; CD339;
Entrez Gene ID	16449
Protein Refseq	NP_038850
UniProt ID	Q63722
Chromosome Location	2 F3; 2 67.73 cM
Pathway	Activated NOTCH1 Transmits Signal to the Nucleus; Constitutive Signaling by NOTCH1 HD Domain Mutants; Constitutive Signaling by NOTCH1 HD+PEST Domain Mutants; Constitutive Signaling by NOTCH1 PEST Domain Mutants; Constitutive Signaling by NOTCH1 t(7;9)(NOTCH1:M1580_K2555) Translocation Mutant; Delta-Notch Signaling Pathway; Disease; FBXW7 Mutants and NOTCH1 in Cancer;
Function	Notch binding; calcium ion binding; protein binding;