



Mouse Anti-Human Collagen Type IV monoclonal antibody, clone JID660 (CABT- L2976)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	This antibody is intended for qualified laboratories to qualitatively identify by light microscopy the presence of associated antigens in sections of formalin-fixed, paraffin-embedded tissue sections using IHC test methods.
Specificity	Human Collagen Type IV
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Clone	JID660
Conjugate	Unconjugated
Applications	IHC
Reconstitution	The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. The concentrated antibody requires dilution in the optimized buffer, to the recommended working dilution range.
Positive Control	Lung, Muscle
Format	Liquid
Size	Predilut: 7 ml, Concentrate: 100 µl, Concentrate: 1 ml

Buffer	Predilute: Antibody Diluent Buffer Concentrate: Tris Buffer, pH 7.3 - 7.7, with 1% BSA
Preservative	< 0.1% Sodium Azide
Storage	Store at 2-8°C. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction	Collagen Type IV is a primary component in the basal lamina that is used as a marker to observe the presence of the lamina and examine its structure. In addition to the epithelial basal lamina, Anti-Collagen Type IV stains mesenchymal components. It is useful for identifying soft tissue cancers including schwannomas and leiomyomas. Anti-Collagen Type IV frequently reacts with these tissues after becoming well-differentiated and malignant. The use of Anti-Collagen Type IV produces more reliable results than non-specific silver reticulum stains when investigating the vascular elements of neoplasms, hemangiopericytoma, angiosarcoma and epithelioid hemangioendothelioma.
Keywords	Arresten;Canstatin;CO4A1_HUMAN;COL4A1;COL4A1 NC1 domain;COL4A2;COL4A3;COL4A4;COL4A5;Collagen Alpha 1(IV) Chain