



Mouse Anti-Human CEA monoclonal antibody, clone JID654 (CABT-L2986)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	This antibody is intended for qualified laboratories to qualitatively identify by light microscopy the presence of associated antigens in sections of formalin-fixed, paraffin-embedded tissue sections using IHC test methods.
Specificity	Human CEA
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Clone	JID654
Conjugate	Unconjugated
Applications	IHC
Reconstitution	The prediluted antibody does not require any mixing, dilution, reconstitution, or titration; the antibody is ready-to-use and optimized for staining. The concentrated antibody requires dilution in the optimized buffer, to the recommended working dilution range.
Positive Control	Colon Adenocarcinoma, Colon Mucosa
Format	Liquid
Size	Predilut: 7 ml, Concentrate: 100 µl, Concentrate: 1 ml
Buffer	Predilute: Antibody Diluent Buffer

Concentrate: Tris Buffer, pH 7.3 - 7.7, with 1% BSA

Preservative	< 0.1% Sodium Azide
---------------------	---------------------

Storage	Store at 2-8°C. Do not freeze.
----------------	--------------------------------

Ship	Wet ice
-------------	---------

BACKGROUND

Introduction

Carcinoembryonic antigen (CEA) describes a set of glycoposphatidyl inositol and transmembrane cell-surface-anchored glycoproteins involved in cell adhesion, differentiation, anoikis, polarization, and tissue architecture. CEA staining, along with Calretinin, CK 5/6, D2-40, HBME-1, Napsin A, MOC-31, and Ber-EP4, is used to help differentiate between adenocarcinoma and mesothelioma. Staining with Anti-CEA is also suggested to be useful in identifying the origin of metastatic adenocarcinoma. CEA is an effective marker for adenocarcinomas of the lung, colon, stomach, esophagus, pancreas, gallbladder, urachus, salivary gland, ovary, and endocervix.

Keywords	CEA;Carcinoembryonic antigen
-----------------	------------------------------
