



# Sheep anti Rabbit Fibrinogen polyclonal antibody (CABT-L394)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	This antibody is specific for rabbit Fg as demonstrated by immunoelectrophoresis and ELISA.
<b>Target</b>	Fibrinogen
<b>Immunogen</b>	Rabbit fibrinogen purified from plasma.
<b>Isotype</b>	IgG
<b>Source/Host</b>	Sheep
<b>Species Reactivity</b>	Rabbit
<b>Purification</b>	Affinity purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	IEP, ELISA
<b>Format</b>	Liquid
<b>Size</b>	0.5 mg
<b>Buffer</b>	10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.
<b>Preservative</b>	None
<b>Storage</b>	Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

## BACKGROUND

## Introduction

Fibrinogen is an abundant plasma protein (5-10  $\mu$ M) produced in the liver. The intact protein has a molecular weight of 340 kDa and is composed of 3 pairs of disulphide-bound polypeptide chains named A $\alpha$ , B $\beta$  and  $\gamma$ . Fibrinogen is a triglobular protein consisting of a central E domain and terminal D domains. Proteolysis by thrombin results in release of Fibrinopeptide A (FPA, A  $\alpha$ 1-16) followed by Fibrinopeptide B (FPB, B $\beta$ 1-14) and the fibrin monomers that result polymerize in a half-overlap fashion to form insoluble fibrin fibrils. The chains of fibrin are referred to as  $\alpha$ ,  $\beta$  and  $\gamma$ , due to the removal of FPA and FPB. The polymerised fibrin is subsequently stabilized by the transglutaminase activated Factor XIII that forms amide linkages between  $\gamma$  chains and, to a lesser extent,  $\alpha$  chains of the fibrin molecules. Proteolysis of fibrinogen by plasmin initially liberates Cterminal residues from the A $\alpha$  chain to produce fragment X (intact DE-D, which is still clottable). Fragment X is further degraded to nonclottable fragments Y (D-E) and D. Fragment Y can be digested into its constituent D and E fragments. Digestion of non-crosslinked fibrin with plasmin is very similar to the digestion of fibrinogen, which results in production of fragments D and E. Degradation of crosslinked fibrin by plasmin results in fragment DD (D-Dimer consisting of the D domains of 2 fibrin molecules crosslinked via the  $\gamma$  chains), fragment E (central E domain) as well as DDE in which fragment E is non-covalently associated with DD. For human crosslinked fibrin, the relative weights of the cleavage fragments produced are: 184 kDa for fragment DD, 92 kDa for D, 50 kDa for E, 1.54 kDa for FP A and 1.57 kDa for FPB.

---

## Keywords

fibrinogen alpha chain;FGA;Fib2;fibrinogen, A alpha polypeptide;

---

# GENE INFORMATION

## Entrez Gene ID

[100353616](#)

## UniProt ID

[G1T0X2](#)

---