



Mouse Anti-Mouse MHC Class I (H-2Kb) Monoclonal antibody, clone AF6-88.5.5.3 (CABT-L4417)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	The AF6-88.5.5.3 monoclonal antibody reacts with the mouse H-2Kb MHC class I alloantigen. MHC class I antigens are heterodimers consisting of one alpha chain (44 kDa) associated with β 2 microglobulin (11.5 kDa). The antigen is expressed by all nucleated cells at varying levels. MHC Class I molecules present endogenously synthesized antigenic peptides to CD8 T cells.
Target	Mouse MHC Class I (H-2Kb)
Immunogen	C57BL/6 mouse splenocytes
Isotype	IgG2a, κ
Source/Host	Mouse
Species Reactivity	Mouse
Clone	AF6-88.5.5.3
Purification	Protein G purified. Purity>95%. Determined by SDS-PAGE
Conjugate	Functional Grade
Applications	in vivo administration, FC
Molecular Weight	150 kDa
Format	0.2 μ M filtered liquid. Purified from tissue culture supernatant in an animal free facility
Concentration	Lot specific

Size	5 mg
Buffer	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free] Endotoxin level: <2EU/mg (<0.002EU/μg). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
Preservative	None
Storage	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction	MHC class I molecules are one of two primary classes of major histocompatibility complex (MHC) molecules (the other being MHC class II) and are found on the cell surface of all nucleated cells in the body.
Keywords	A 28;A 9;Antigen presenting molecule;Aw 24;Aw 68;CLASS I HISTOCOMPATIBILITY ANTIGEN;H2 K1;H2K;HLA A;HLA class I histocompatibility antigen A 1 alpha chain

GENE INFORMATION

Official Symbol	MHC Class I
Synonyms	A 28; A 9; Antigen presenting molecule; Aw 24; Aw 68; CLASS I HISTOCOMPATIBILITY ANTIGEN; H2 K1; H2K; HLA A; HLA class I histocompatibility antigen A 1 alpha chain
References	Paschall, A. V., et al. (2015). "IFN regulatory factor 8 represses GM-CSF expression in T cells to affect myeloid cell lineage differentiation." J Immunol 194(5): 2369-2379. PubMed;