



Mouse Anti-Mouse CTLA-4 (CD152) Monoclonal antibody, clone 9D9 (CABT-L4442)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview

The 9D9 monoclonal antibody reacts with mouse CTLA-4 (cytotoxic T lymphocyte antigen-4) also known as CD152. CTLA-4 is a 33 kDa cell surface receptor encoded by the *Ctla4* gene that belongs to the CD28 family of the Ig superfamily. CTLA-4 is expressed on activated T and B lymphocytes. CTLA-4 is structurally similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to the B7 family members B7-1 (CD80) and B7-2 (CD86). Upon ligand binding, CTLA-4 negatively regulates cell-mediated immune responses. CTLA-4 plays roles in induction and/or maintenance of immunological tolerance, thymocyte development, and regulation of protective immunity. The critical role of CTLA-4 in immune down-regulation has been demonstrated in CTLA-4 deficient mice, which succumb at 3-5 weeks of age due to the development of a lymphoproliferative disease. CTLA-4 is among a group of inhibitory receptors being explored as cancer treatment targets through immune checkpoint blockade.

Target	Mouse CTLA-4 (CD152)
Immunogen	CHO cell line transfected with mouse CD80
Isotype	IgG2b
Source/Host	Mouse
Species Reactivity	Mouse
Clone	9D9
Purification	Protein A purified. Purity>95%. Determined by SDS-PAGE
Conjugate	Functional Grade
Applications	in vivo CTLA-4 neutralization, WB

Molecular Weight	150 kDa
Format	0.2 µM filtered liquid. Purified from tissue culture supernatant in an animal free facility
Concentration	Lot specific
Size	5 mg
Buffer	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free] Endotoxin level: <2EU/mg (<0.002EU/µg). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
Preservative	None
Storage	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hashimoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases.

Keywords CTLA-4

GENE INFORMATION

Official Symbol cytotoxic T-lymphocyte-associated protein 4

Synonyms CTLA-4

References Dai, M., et al. (2015). "Curing mice with large tumors by locally delivering combinations of immunomodulatory antibodies." Clin Cancer Res 21(5): 1127-1138. PubMed;Balachandran, V. P., et al. (2011). "Imatinib potentiates antitumor T cell responses in gastrointestinal stromal tumor through the inhibition of Ido." Nat Med 17(9): 1094-1100. PubMed;Curran, M. A., et al. (2011). "Combination CTLA-4 blockade and 4-1BB activation enhances tumor rejection by

increasing T-cell infiltration, proliferation, and cytokine production." PLoS One 6(4): e19499.
PubMed;
