



Mouse Anti-Human IFN γ Monoclonal antibody, clone B133.5 (CABT-L4487)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview

The B133.5 monoclonal antibody reacts with human IFN γ (interferon gamma) a 20 kDa soluble pleiotropic cytokine and the sole member of the type II class of interferons. IFN γ is primarily produced by activated lymphocytes including T, B, NK cells, and ILCs. IFN γ exerts immunoregulatory, anti-proliferative, anti-viral, and proinflammatory activities and plays an important role in activation, growth, and differentiation of T and B lymphocytes, macrophages, NK cells and other non-hematopoietic cell types. Additionally, IFN γ induces the production of cytokines, Fc receptor, and adhesion molecules and up-regulates MHC class I and II antigen expression by antigen presenting cells during an immune response. IFN γ has also been shown to modulate macrophage effector functions, influence isotype switching and induce the secretion of immunoglobulins by B cells. IFN γ signals through the IFN gamma receptor which exists as a heterodimer composed of CD119 (IFN gamma receptor 1) and AF-1 (IFN gamma receptor 2). The IFN γ receptor is expressed ubiquitously on almost all cell types with the exception of mature erythrocytes. The B133.5 antibody is a neutralizing antibody.

Target	Human IFN γ
Immunogen	Recombinant human IFN γ
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	B133.5
Purification	Protein A purified. Purity>95%. Determined by SDS-PAGE
Conjugate	Functional Grade

Applications	in vitro IFN γ neutralization
Molecular Weight	150 kDa
Format	0.2 μ M filtered liquid. Purified from tissue culture supernatant in an animal free facility
Concentration	Lot specific
Size	5 mg
Buffer	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free] Endotoxin level: <2EU/mg (<0.002EU/ μ g). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
Preservative	None
Storage	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction	This gene encodes a member of the type II interferon family. The protein encoded is a soluble cytokine with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. Mutations in this gene are associated with aplastic anemia.
Keywords	IFNG;interferon, gamma;IFG;IFI;interferon gamma;IFN-gamma;immune interferon

GENE INFORMATION

Official Symbol	interferon, gamma
Synonyms	IFNG; interferon, gamma; IFG; IFI; interferon gamma; IFN-gamma; immune interferon
References	Mitson-Salazar, A., et al. (2015). "Hematopoietic prostaglandin D synthase defines a proeosinophilic pathogenic effector human T2 cell subpopulation with enhanced function." J Allergy Clin Immunol. pii: S0091-6749(15)01186-0. PubMed;