



Armenian Hamster Anti-Mouse TNF α Monoclonal antibody, clone TN3-19.12 (CABT-L4492)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	The TN3-19.12 monoclonal antibody reacts with mouse, rat, and rabbit TNF α (tumor necrosis factor-alpha) a multifunctional proinflammatory cytokine
Target	Mouse/Rat/rabbit TNF α
Immunogen	Recombinant mouse TNF α
Isotype	IgG, κ
Source/Host	Armenian Hamster
Species Reactivity	Rat, Mouse, Rabbit
Clone	TN3-19.12
Purification	Protein G purified. Purity>95%. Determined by SDS-PAGE
Conjugate	Functional Grade
Applications	in vivo TNF α neutralization, FC
Molecular Weight	150 kDa
Format	0.2 μ M filtered liquid. Purified from tissue culture supernatant in an animal free facility
Concentration	Lot specific
Size	5 mg

Buffer	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free] Endotoxin level: <2EU/mg (<0.002EU/μg). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
Preservative	None
Storage	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction

The TN3-19.12 monoclonal antibody reacts with mouse, rat, and rabbit TNF α (tumor necrosis factor-alpha) a multifunctional proinflammatory cytokine. TNF α exists as a soluble 17 kDa monomer, which forms homotrimers in circulation or as a 26 kDa membrane-bound form. TNF α belongs to the TNF superfamily of cytokines and signals through its two receptors, TNFR1 and TNFR2 which can be activated by both the soluble trimeric and membrane-bound and forms of TNF α . TNF α is primarily produced by macrophages in response to foreign antigens such as bacteria (lipopolysaccharides), viruses, and parasites as well as mitogens and other cytokines but can also be expressed by monocytes, neutrophils, NK cells, CD4 T cells and some specialized dendritic cells. TNF α is known to play key roles in a wide spectrum of biological processes including immunoregulation, cell proliferation, differentiation, apoptosis, antitumor activity, inflammation, anorexia, cachexia, septic shock, hematopoiesis, and viral replication. TNF α dysregulation has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Mouse and human TNF α share 79% amino acid sequence identity however, mouse TNF α is glycosylated while human TNF α is not. TNF α knockout animals display defects in response to bacterial infection, characterized by defects in forming organized follicular dendritic cell networks and germinal centers with a lack of primary B cell follicles. The TN3-19.12 antibody can neutralize the bioactivity of natural or recombinant TNF- α .

Keywords APC1;APC1 protein;Cachectin;DIF;Differentiation inducing factor;Macrophage cytotoxic factor;MCF;Necrosin;TNF a;TNF alpha

GENE INFORMATION

Official Symbol	Tumor necrosis factor alpha
Synonyms	APC1; APC1 protein; Cachectin; DIF; Differentiation inducing factor; Macrophage cytotoxic factor; MCF; Necrosin; TNF a; TNF alpha
References	Xiong, H., et al. (2016). "Innate Lymphocyte/Ly6C Monocyte Crosstalk Promotes Klebsiella

