

Human Anti-SARS-CoV-2 Spike RBD Monoclonal antibody, clone 1P2P3P (CABT-L4572)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	SARS-CoV-2 Spike RBD
Isotype	IgG
Source/Host	Human
Species Reactivity	SARS-CoV-2
Clone	1P2P3P
Purification	Affinity purified
Conjugate	Unconjugated
Applications	Suitable for assay development and WB Cocktail of anti-SARS-CoV-2 spike glycoprotein monoclonal antibody, clone 1P,2P, and 3P. Clone 1P, 2P, 3P are all available upon request.
Format	Liquid
Concentration	Lot specific
Size	100 μg, 1 mg
Buffer	PBS,ph7.4
Preservative	None
Storage	Store -20°C for long term.

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

BACKGROUND

Introduction

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigenrelated cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Keywords

SARS-CoV-2 spike glycoprotein;SARS-CoV-2 spike;SARS-CoV-2;SARS-CoV;2019nCoV;Coronavirus;Human Coronavirus;HCoV;SARS;SARS CoV;SARS-CoV-2 S1;2019-nCoV S1;COVID-19;SARS-CoV-2 S1