



Goat anti Human Plasminogen polyclonal antibody [HRP] (CABT-L479)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	Prior to conjugation, this antibody was specific for Pg as demonstrated by immunoelectrophoresis and ELISA.
Target	Plasminogen
Immunogen	Human plasminogen purified from plasma.
Isotype	IgG
Source/Host	Goat
Species Reactivity	Human
Purification	Affinity purified
Conjugate	HRP
Applications	IEP, ELISA
Format	Liquid
Size	100 µg
Buffer	A buffered stabilizer solution containing 50% (v/v) glycerol.
Preservative	None
Storage	Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use. Avoid exposure to sodium azide as this is an inhibitor of peroxidase activity.

BACKGROUND

Introduction	Plasminogen (Pg) is synthesized in the liver and circulates in plasma at a concentration of ~200 µg/ml (~2.3 µM). Plasminogen is a single-chain glycoprotein of ~88 kDa that consists of a catalytic domain followed by five kringle structures. Within these kringle structures are four low-affinity lysine binding sites and one high-affinity lysine binding site. It is through these lysine binding sites that plasminogen binds to fibrin and to α2 Antiplasmin. Native plasminogen (glu-plasminogen) exists in two variants that differ in their extent of glycosylation, and each variant has up to six isoelectric forms with respect to sialic acid content, for a total of 12 molecular forms. Activation of glu-plasminogen by the plasminogen activators urokinase (UPA), or tissue plasminogen activator (tPA) occurs by cleavage after residue Arg560 to produce the two-chain active serine protease plasmin. In a positive feedback reaction, the plasmin generated cleaves an ~8 kDa peptide from glu-plasminogen, producing lys77-plasminogen which has a higher affinity for fibrin and when bound is a preferred substrate for plasminogen activators such as urokinase. Additional activators of plasminogen include kallikrein and activated factor XII. The primary inhibitor of plasmin in plasma is α2 Antiplasmin. Other physiological inhibitors of plasmin include α2 macroglobulin and antithrombin.
--------------	---

Keywords	PLG;plasminogen;plasmin;
----------	--------------------------

GENE INFORMATION

Entrez Gene ID	5340
UniProt ID	P00747