



Anti-CCL17 polyclonal antibody (CPBT-65034RH)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Rabbit anti Human CCL17 antibody specifically recognizes human CCL17, otherwise known as TARC (thymus and activation-regulated chemokine), a member of the intercrine beta (chemokine CC) family, predominantly produced by thymic dendritic cells, which specifically acts as a chemotactic factor for T cells, and signals through binding to the CCR4 receptor. ELISA This antibody may be used in an indirect ELISA or as a capture reagent in a sandwich ELISA with B as the detection antibody.
Specificity	CCL17
Immunogen	Recombinant human CCL17
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Conjugate	Unconjugated
Applications	ELISA; FA; WB
Format	Purified IgG - lyophilised
Size	100 µg
Preservative	None
Storage	in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

GENE INFORMATION

Gene Name	CCL17 chemokine (C-C motif) ligand 17 [Homo sapiens (human)]
Official Symbol	CCL17
Synonyms	CCL17; chemokine (C-C motif) ligand 17; TARC; ABCD-2; SCYA17; A-152E5.3; C-C motif chemokine 17; CC chemokine TARC; T cell-directed CC chemokine; small-inducible cytokine A17; thymus and activation-regulated chemokine; small inducible cytokine subfamily A
Entrez Gene ID	6361
Protein Refseq	NP_002978
UniProt ID	Q92583
Chromosome Location	16q13
Pathway	Chemokine receptors bind chemokines; Chemokine signaling pathway; Class A/1 (Rhodopsin-like receptors); Cytokine-cytokine receptor interaction; Defective ACTH causes Obesity and Pro-opiomelanocortinin deficiency (POMCD); Disease; GPCR ligand binding; IL4-mediated signaling events;
Function	CCR4 chemokine receptor binding; chemokine activity; receptor binding;