



Inactivated Anatid alphaherpesvirus 1 (AnHV-1) (DAG-WT1134)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Conjugate | N/A |
|---------------|--|
| Applications | Immunoassays |
| Format | Liquid |
| Concentration | Batch dependent - please inquire should you have specific requirements |
| Size | 1 mL |
| Preservative | None |
| Storage | Store at -20°C |

BACKGROUND

| Keywords | is not zoonotic. Duck plague; Anatid alphaherpesvirus 1 |
|--------------|--|
| | factor in the spread of this disease as they are often asymptomatic carriers of disease. The incubation period is three to seven days. Birds as young as one week old can be infected. DEV |
| | horizontally—through contaminated water and direct contact. Migratory waterfowl are a major |
| | mortality rates in flocks of ducks, geese, and swans. It is spread both vertically and |
| Introduction | Duck plague (also known as duck viral enteritis) is a worldwide disease caused by Anatid alphaherpesvirus 1 (AnHV-1) of the family Herpesviridae that causes acute disease with high |

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221