

## Sheep Anti Bovine IgG2 Polyclonal Antibody, AP

DPBT-67026SB Sheep(IgG2) Lot. No. (See product label)

## PRODUCT INFORMATION

**Product Overview** Sheep Anti Bovine IgG2,AP

*Immunogen* Purified bovine IgG2

*Host* Sheep

Isotype Polyclonal IgG

SpeciesBovineConjugationAPApplicationsELISA,

**Dilution** ELISA: 1/1000 - 1/10000

## **PACKAGING**

Format Purified IgG conjugated to Alkaline Phosphatase - liquid

**Protein Concentration** IgG concentration 0.5mg/ml

Buffer 50mM HEPES, 0.1M NaCl, 1mM MgCl2, 0.1mM ZnCl2

Storage Store at +4 °C.DO NOT FREEZE. This product should be stored undiluted. Should this product contain

a precipitate we recommend microcentrifugation before use.

Preservative 0.09% Sodium Azide (NaN3)

Shelf Life 12 months from date of despatch.

## **BACKGROUND**

Immunoglobulin G (IgG) are antibody molecules. Each IgG is composed of four peptide chains - two

heavy chains γ and two light chains. Each IgG has two antigen binding sites. Other Immunoglobulins may be described in terms of polymers with the IgG structure considered the monomer. IgG molecules are synthesized and secreted by plasma B cells. IgG antibodies are large molecules of about 150 kDa composed of 4 peptide chains. It contains 2 identical heavy chains of about60kDa and 2 identical light chains of about 25 kDa, thus a tetrameric quaternary structure. The two heavy chains are linked to each other and to a light chain each by disulfide bonds. The resulting tetramer has two identical halves, which together form the Y-like shape. Each end of the fork contains an identical antigen binding site. The Fc regions of IgGs bear a highly conserved N-glycosylation site. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α-2,6-linked sialic acid

residues.

Keywords Ig gamma-2 chain C region antibody; immunoglobulin Gm2; immunoglobulin heavy constant gamma 2

(G2m marker); immunoglobulin heavy constant, gamma 2; IgG2; Immunoglobulin G2; IgG2γ; Immunoglobulin G2γ; IgG2 heavy chain, Immunoglobulin G2 heavy chain; IgG2γheavy chain;

Immunoglobulin G2γheavy chain