

Goat Anti Chicken IgG (Fc) Polyclonal Antibody, AP

DPBT-67115GC Goat(IgG) Lot. No. (See product label)

PRODUCT INFORMATION

Product OverviewGoat Anti Chicken IgG (Fc),APImmunogenPurified chicken IgG (Fc) fragment.

Host Goat

Isotype Polyclonal IgG

SpeciesChickenConjugationAPApplicationsELISA,

Dilution ELISA: 1/5000 - 1/50000

PACKAGING

Format Purified IgG conjugated to Alkaline Phosphatase - liquid

Protein Concentration IgG concentration 0.5mg/ml

Buffer 50mM HEPES, 0.1M NaCl, 1mM MgCl2, 0.1mM ZnCl2

Storage Store at +4 °C.DO NOT FREEZE. This product should be stored undiluted. Should this product contain

a precipitate we recommend microcentrifugation before use.

Preservative 0.09%Sodium Azide0.2%Bovine Serum Albumin

Shelf Life 12 months from date of despatch.

BACKGROUND

Immunoglobulin G (IgG) are antibody molecules. Each IgG is composed of four peptide chains - two

heavy chains γ and two light chains. Each IgG has two antigen binding sites. Other Immunoglobulins may be described in terms of polymers with the IgG structure considered the monomer. IgG molecules are synthesized and secreted by plasma B cells. IgG antibodies are large molecules of about 150 kDa composed of 4 peptide chains. It contains 2 identical heavy chains of about60kDa and 2 identical light chains of about 25 kDa, thus a tetrameric quaternary structure. The two heavy chains are linked to each other and to a light chain each by disulfide bonds. The resulting tetramer has two identical halves, which together form the Y-like shape. Each end of the fork contains an identical antigen binding site. The Fc regions of IgGs bear a highly conserved N-glycosylation site. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α-2,6-linked sialic acid

residues.

Keywords Ig gamma 1 chain C region; IGHG1; Immunoglobin heavy constant gamma 1; Immunoglobulin G; IgG;

IgG heavy chain; Immunoglobulin G heavy chain