

Rabbit Anti Sheep IgA Polyclonal Antibody, FITC

DPBT-67213RS Rabbit(IgA)

Lot. No. (See product label)

PRODUCT INFORMATION

Product Overview	Rabbit Anti Sheep IgA,FITC
Host	Rabbit
Isotype	Polyclonal IgG
Species	Sheep
Conjugation	FITC
Applications	IHC, ELISA, FCM, IP, WB
Dilution	FCM: 1/200 - 1/2000

PACKAGING

Format	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid
Protein Concentration	IgG concentration 1.0 mg/ml
Buffer	Phosphate buffered saline
Storage	Store at +4 °C or at -20 °C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Preservative	0.09% Sodium Azide 0.2% Bovine Serum Albumin
Shelf Life	18 months from date of despatch.

BACKGROUND

Introduction	Immunoglobulin A (IgA) is an antibody that plays a critical role in mucosal immunity. More IgA is produced in mucosal linings than all other types of antibody combined; between three and five grams are secreted into the intestinal lumen each day. This accumulates to 75% of the total immunoglobulin produced in the entire body. IgA has two subclasses (IgA1 and IgA2) and can exist in a dimeric form called secretory IgA (sIgA). In its secretory form, IgA is the main immunoglobulin found in mucous secretions, including tears, saliva, colostrum and secretions from the genitourinary tract, gastrointestinal tract, prostate and respiratory epithelium. It is also found in small amounts in blood. The secretory component of sIgA protects the immunoglobulin from being degraded by proteolytic enzymes, thus sIgA can survive in the harsh gastrointestinal tract environment and provide protection against microbes that multiply in body secretions. IgA is a poor activator of the complement system, and opsonises only weakly. Its heavy chains are of the type α .
Keywords	Ig alpha 1 chain C region; Ig alpha 2 chain C region; IGHA1; IGHA2; Immunoglobulin heavy constant alpha 1; Immunoglobulin heavy constant alpha 2 A2m marker; Immunoglobulin heavy constant alpha 2; IgA; Immunoglobulin A; IgA α ; Immunoglobulin A α ; IgA heavy chain, Immunoglobulin A heavy chain; IgA heavy chain; Immunoglobulin A heavy chain