








Recombinant Porphyromonas gingivalis Gingipain R1(rgpA),partial

Product Code CSB-MP338957PQP Relevance Thrombin-like snake venom serine protease that acts as an anticoagulant. It cleaves fibrinogen (FGA) to split off the A-fibrinopeptides (A, AY and AP), but not the B-fibrinopeptide. The resulting fibrin polymers are imperfectly formed and much smaller in size (1 to 2 um long) than the fibrin polymers produced by the action of thrombin. These ancrod-induced microthrombia re friable, unstable,		
cleaves fibrinogen (FGA) to split off the A-fibrinopeptides (A, AY and AP), but not the B-fibrinopeptide. The resulting fibrin polymers are imperfectly formed and much smaller in size (1 to 2 um long) than the fibrin polymers produced by the action of thrombin. These ancrod-induced microthrombi are friable, unstable, urea-soluble and have significantly degraded alpha chains. They do not crosslink to form thrombi. They are markedly susceptible to digestion by plasmin and are rapidly roved from circulation by either reticuloendothelial phagocytosis or normal fibrinolysis, or both. Anticoagulation through the roval of fibrinogen from the blood is rapid, occurring within hours following its administration. It does not activate plasminogen and does not degrade preformed, fully cross-linked thrombin fibrin. It also reduces the level of plasminogen activator (PLAT) from the endothelium. The profibrinolytic effect of these 2 actions appears to be limited to local microthrombus degradation. Storage The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. Uniprot No. P28784 Storage Buffer Tris-based buffer,50% glycerol Product Type Recombinant Protein Species Porphyromonas gingivalis Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIKICYDPGVTPKNIDAFNGGISLVNYTGIGSETA WGTSHEGTTHVKQLTINSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGGVTMNGMFAM VEKYKKDGEKMLDTWYFGPSSLVATLTGNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others	Product Code	CSB-MP338957PQP
storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. Uniprot No. P28784 Storage Buffer Tris-based buffer,50% glycerol Product Type Recombinant Protein Species Porphyromonas gingivalis Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNAGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Mammalian cell	Relevance	cleaves fibrinogen (FGA) to split off the A-fibrinopeptides (A, AY and AP), but not the B-fibrinopeptide. The resulting fibrin polymers are imperfectly formed and much smaller in size (1 to 2 um long) than the fibrin polymers produced by the action of thrombin. These ancrod-induced microthrombi are friable, unstable, urea-soluble and have significantly degraded alpha chains. They do not cross-link to form thrombi. They are markedly susceptible to digestion by plasmin and are rapidly roved from circulation by either reticuloendothelial phagocytosis or normal fibrinolysis, or both. Anticoagulation through the roval of fibrinogen from the blood is rapid, occurring within hours following its administration. It does not activate plasminogen and does not degrade preformed, fully cross-linked thrombin fibrin. It also reduces the level of plasminogen activator inhibitor (PAI) and may stimulate the release of tissue plasminogen activator (PLAT) from the endothelium. The profibrinolytic effect of these 2 actions appears to be limited to
Storage Buffer Tris-based buffer,50% glycerol Product Type Recombinant Protein Species Porphyromonas gingivalis Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKIPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Source Mammalian cell	Storage	storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12
Product Type Recombinant Protein Species Porphyromonas gingivalis Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Source Mammalian cell	Uniprot No.	P28784
Species Porphyromonas gingivalis Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Source Mammalian cell	Storage Buffer	Tris-based buffer,50% glycerol
Purity Greater than 90% as determined by SDS-PAGE. Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Mammalian cell	Product Type	Recombinant Protein
Sequence YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTAN AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Mammalian cell	Species	Porphyromonas gingivalis
AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS VSDAPELLRS Research Area Others Mammalian cell	Purity	Greater than 90% as determined by SDS-PAGE.
Source Mammalian cell	Sequence	AIQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNE VFIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGE SDIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGK PTGTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAM VEKYKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCD YNGAIATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINT NGEPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLS
	Research Area	Others
Gene Names rgpA	Source	Mammalian cell
	Gene Names	rgpA



CUSABIO TECHNOLOGY LLC








Expression Region	228-720aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	58.0kDa
Protein Description	Partial

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.