

Recombinant Human High affinity immunoglobulin epsilon receptor subunit alpha(FCER1A),partial

Product Code	CSB-EP008532HU
Relevance	Binds to the Fc region of immunoglobulins epsilon. High affinity receptor. Responsible for initiating the allergic response. Binding of allergen to receptor- bound IgE leads to cell activation and the release of mediators (such as histamine) responsible for the manifestations of allergy. The same receptor also induces the secretion of important lymphokines.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P12319
Storage Buffer	Tris-based buffer,50% glycerol
Alias	c-epsilon RI-alpha Short name: FcERI IgE Fc receptor subunit alpha
Product Type	Recombinant Protein
Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	VPQKPKVSLNPPWNRIFKGENVTLTCNGNNFFEVSSTKWFHNGSLSEETNSS LNIVNAKFEDSGEYKCQHQQVNESEPVYLEVFSDWLLLQASAEVVMEGQPLF LRCHGWRNWDVYKVIYYKDGEALKYWYENHNISITNATVEDSGTYYCTGKVW QLDYESEPLNITVIKAPREKYWLQ
Lead Time	3-7 business days
Research Area	Immunology
Source	E.coli
Gene Names	FCER1A
Protein Names	Recommended name: High affinity immunoglobulin epsilon receptor subunit alpha Alternative name(s): Fc-epsilon RI-alpha Short name= FcERI IgE Fc receptor subunit alpha
Expression Region	26-205aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-SUMO-tagged
Mol. Weight	37.0kDa
Protein Description	Extracellular Domain
Image	

1



CUSABIO TECHNOLOGY LLC

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🧉

kDa	М	
116.0	-	
66.2		
45.0		
35.0		-
25.0		
18.4		
14.4		

(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.