




Recombinant Human Nephrocystin-1(NPHP1),partial

Product Code	CSB-EP015985HU
Relevance	Together with BCAR1 it may play a role in the control of epithelial cell polarity. Involved in the organization of apical junctions in kidney cells together with NPHP4 and RGPRI1L/NPHP8 . Does not se to be strictly required for ciliogenesis . Ses to help to recruit PTK2B/PYK2 to cell matrix adhesions, thereby initiating phosphorylation of PTK2B/PYK2 and PTK2B/PYK2-dependent signaling. May play a role in the regulation of intraflagellar transport (IFT) during cilia assbly. Required for normal retina development. In connecting photoreceptor cilia influences the movent of some IFT proteins such as IFT88 and WDR19. Involved in spermatogenesis .
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	O15259
Storage Buffer	Tris-based buffer,50% glycerol
Alias	Juvenile nephronophthisis 1 protein
Product Type	Recombinant Protein
Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MLARRQRDPLQALRRRNQELKQQVDSLLSESQLEALEPNKRQHIIYQRCIQLK QAIDENKNALQKLSKADESAPVANYNQRKEEEHTLLDKLTQQQLQGLAVTISRE NIT
Research Area	Metabolism
Source	E.coli
Gene Names	NPHP1
Expression Region	1-109aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal GST-tagged
Mol. Weight	39.7kDa
Protein Description	Partial
Image	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.