




Recombinant Human Perforin-1 (PRF1)

Product Code	CSB-EP018668HU
Relevance	Plays a key role in secretory granule-dependent cell death, and in defense against virus-infected or neoplastic cells. Plays an important role in killing other cells that are recognized as non-self by the immune syst, e.g. in transplant rejection or some forms of autoimmune disease. Can insert into the mbrane of target cells in its calcium-bound form, oligomerize and form large pores. Promotes cytolysis and apoptosis of target cells by facilitating the uptake of cytotoxic granzymes.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P14222
Storage Buffer	Tris-based buffer,50% glycerol
Alias	Cytolysin;Lymphocyte pore-forming protein ;PFP
Product Type	Recombinant Protein
Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	PCHTAARSECKRSHKFVPGAWLAGEGVDVTSLRRSGSFPVDTQRFLRPDGT CTLCENALQEGTLQRLPLALTNWRAQSGGCQRHVTRAKVSSTEAVARDAARS IRNDWKVGLDVTPKPTSNVHVSVAGSHSQAANFAAQKTHQDQYSFSTDTVEC RFYSFHVVHTPPLHPDFKRALGDLPHHFNASTQPAYLRLISNYGTHFIRAVELG GRISALTALRTCELALEGLTDNEVEDCLTVEAQVNIHIGHSISAEAKACEEKKKK HKMTASFHQTYRERHSEVVGGHHTSINDLLFGIQAGPEQYSAWVNSLPGSPG LVDYTLLEPLHVLLDSQDPRREALRRALSQYLTDRARWRDCSRPCPPGRQKSP RDPCQCVCCHGSAVTTQDCCPRQRGLAQLEVTFIQAWGLWGDWFTATDAYVK LFFGGQELRTSTVWDNNNPIWSVRLDFGDVLLATGGPLRLQVWDQDSGRDD DLLGTCDQAPKSGSHEVRCNLNHGHLKFRYHARCLPHLGGGTCLDYVPQMLL GEPGNGRSGAVW
Lead Time	3-7 business days
Research Area	Immunology
Source	E.coli
Gene Names	PRF1
Expression Region	22-555aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-SUMO-tagged
Mol. Weight	75.2kDa

**Protein Description**

Full Length of Mature Protein

Image

(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.