







TUBB Monoclonal Antibody

| Product Code | CSB-MA171192 |
|----------------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Raised In | Mouse |
| Species Reactivity | Human,Mouse,Rat |
| Specificity | Theβ-tubulin antibody can detects endogenousβ-tubulin protein. |
| Tested Applications | ELISA,WB;Recommended dilution:WB:1:500-1:5000 |
| Relevance | Microtubules are constituent parts of the mitotic apparatus, cilia, flagella, and elements of the cytoskeleton. They consist principally of 2 soluble proteins, alpha- and beta-tubulin, each of about 55,000 Da. Antibodies against beta Tubulin are useful as loading controls for Western Blotting. However it should be noted that levels of β -Tubulin may not be stable in certain cells. For example, expression of β -Tubulin in adipose tissue is very low and therefore β -Tubulin should not be used as loading control for these tissues. |
| Form | Mouse IgG1 in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Purification Method | Antigen Affinity Purified |
| Clonality | Monoclonal |
| Alias | beta 5-tubulin; beta Ib tubulin; beta-4 tubulin; M40; MGC117247; MGC16435; |
| | OK/SW-cl.56; TBB5; TUBB; TUBB1; TUBB5 |