






AP1S3 Antibody

868EA01HU
t, store at -20°C or -80°C. Avoid repeated freeze.
t Human AP-1 complex subunit sigma-3 protein (1-104AA)
Recommended dilution: WB:1:1000-1:5000
athrin-associated adaptor protein complex 1 that plays a role in ng in the late-Golgi/trans-Golgi network (TGN) and/or endosomes. plexes mediate both the recruitment of clathrin to membranes and on of sorting signals within the cytosolic tails of transmembrane ules.
ated
e: 0.03% Proclin 300 e: 50% Glycerol, 0.01M PBS, PH 7.4
in G purified
ex subunit sigma-3 (Adaptor protein complex AP-1 subunit Adaptor-related protein complex 1 subunit sigma-1C) (Clathrin otein complex 1 sigma-1C small chain) (Golgi adaptor HA1/AP1 na-1C subunit) (Sigma 1C subunit of AP-1 clathrin) (Sigma-adaptin C-adaptin), AP1S3
Western blot



20kDa

All lanes: AP1S3 antibody at 2µg/ml + A549

whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 19, 20, 13 kDa Observed band size: 19 kDa



CUSABIO TECHNOLOGY LLC



