


Image





CHRM3 Antibody

Product Code	CSB-PA005383LA01HU
Abbreviation	Muscarinic acetylcholine receptor M3
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P20309
Immunogen	Recombinant Human Muscarinic acetylcholine receptor M3 protein (253-492AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200
Relevance	The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is Pi turnover.
Form	Liquid
Form Conjugate	Liquid Non-conjugated
Conjugate	Non-conjugated Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Muscarinic acetylcholine receptor M3, CHRM3
Conjugate Storage Buffer Purification Method Isotype Clonality Alias Species	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Muscarinic acetylcholine receptor M3, CHRM3 Human



IHC image of CSB-PA005383LA01HU diluted at 1:400 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized tissue using an HRP conjugated SP system.




CUSABIO TECHNOLOGY LLC











Immunofluorescence staining of PC-3 cells with CSB-PA005383LA01HU at 1:133, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized tissue using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C.The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG (H+L).