


**Image** 





## KRT18 Antibody

Product Code	CSB-PA005823
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P05783
Immunogen	Synthesized peptide derived from Human Keratin 18.
Raised In	Rabbit
Species Reactivity	Human
Specificity	The antibody detects endogenous levels of total Keratin 18 protein.
<b>Tested Applications</b>	ELISA,WB,IHC,IF;WB:1:500-1:3000,IHC:1:50-1:100,IF:1:100-1:500
Relevance	Involved in the uptake of thrombin-antithrombin complexes by hepatic cells By similarity. When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.
	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul
Form	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441.
Form Purification Method	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,
	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  The antibody was affinity-purified from rabbit antiserum by affinity-
Purification Method	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Purification Method Clonality	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Polyclonal
Purification Method  Clonality  Alias	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Polyclonal  Keratin type I cytoskeletal 18; Cytokeratin-18; CK-18 K18;
Purification Method  Clonality  Alias  Product Type	Thomas M. Magin, J. Cell Biol., Mar 1998; 140: 1441. Carlos Caul  Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Polyclonal  Keratin type I cytoskeletal 18; Cytokeratin-18; CK-18 K18;  Polyclonal Antibody



Immunohistochemical analysis of paraffinembedded human lung carcinoma tissue using Keratin 18 antibody.
















Immunofluorescence analysis of HeLa cells, using Keratin 18 antibody.



Western blot analysis of extracts from HeLa cells, using Keratin 18 antibody.



Western blot analysis of extracts from 293 cells (Lane 2), CoLo cells (Lane 3) and HepG2 cells (Lane 4), using Keratin 18 antiobdy. The lane on the left is treated with systhesized peptide.