


DLK1 Antibody

Product Code	CSB-PA006945LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P80370
Immunogen	Recombinant Human Protein delta homolog 1 protein (24-303AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200, IF:1:50-1:200
Relevance	May have a role in neuroendocrine differentiation.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Protein delta homolog 1 (DLK-1) (pG2) [Cleaved into: Fetal antigen 1 (FA1)], DLK1, DLK
Species	Human
Research Area	Signal Transduction
Target Names	DLK1

Image



Western blot


All lanes: protein delta homolog 1 antibody at 2μg/ml + U251 whole cell lysate

Secondary


Goat polyclonal to Rabbit IgG at 1/10000 dilution

Predicted band size: 42, 34 kDa


Observed band size: 42 kDa



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA006945LA01HU at dilution of 1:100



Immunofluorescent analysis of A549 cells using CSB-PA006945LA01HU at dilution of 1:100 and Cy3-conjugated Goat Anti-Rabbit IgG



Western Blot

Positive WB detected in: A549 whole cell lysate, MCF-7 whole cell lysate, HepG2 whole cell lysate, SH-SY5Y whole cell lysate, Rat brain tissue


All lanes: DLK1 antibody at 4.6 µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 42, 34 kDa

Observed band size: 42 kDa



IHC image of CSB-PA006945LA01HU diluted at 1:300 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HepG2 cells with CSB-PA006945LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).