





ENOX1 Antibody, HRP conjugated

Product Code CSB-PA007676LB01HU Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q8TC92 Immunogen Recombinant Human Ecto-NOX disulfide-thiol exchanger 1 protein (101-340AA) Raised In Rabbit Species Reactivity Human Tested Applications ELISA Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constitutents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (CNOX) (Includes: Hydroquinone [NADH] oxidase) (cCNOX) (cNOX) (Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)		
Uniprot No. Q8TC92 Immunogen Recombinant Human Ecto-NOX disulfide-thiol exchanger 1 protein (101-340AA) Raised In Rabbit Species Reactivity Human Tested Applications ELISA Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (CPIOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1); ENOX1 Species Homo sapiens (Human) Research Area Cancer	Product Code	CSB-PA007676LB01HU
Raised In Rabbit Species Reactivity Human Tested Applications Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method softype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit Species Reactivity Human Tested Applications ELISA Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Uniprot No.	Q8TC92
Species Reactivity Human Tested Applications ELISA Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1); Prot	Immunogen	Recombinant Human Ecto-NOX disulfide-thiol exchanger 1 protein (101-340AA)
Tested Applications ELISA Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1); ENOX1 Species Homo sapiens (Human) Research Area Cancer	Raised In	Rabbit
Relevance Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Species Reactivity	Human
NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock. Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Tested Applications	ELISA
Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Relevance	NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Conjugate	HRP
Clonality Polyclonal Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Storage Buffer	
Clonality Polyclonal Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Purification Method	>95%, Protein G purified
Alias Ecto-NOX disulfide-thiol exchanger 1 (Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Isotype	IgG
keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase (EC 1)], ENOX1 Species Homo sapiens (Human) Research Area Cancer	Clonality	Polyclonal
Research Area Cancer	Alias	keeping constitutive hydroquinone [NADH] oxidase) (cCNOX) (Cell proliferation-inducing gene 38 protein) (Constitutive Ecto-NOX) (cNOX) [Includes: Hydroquinone [NADH] oxidase (EC 1); Protein disulfide-thiol oxidoreductase
	Species	Homo sapiens (Human)
Target Names ENOX1	Research Area	Cancer
	Target Names	ENOX1