




ERCC2 Antibody

Product Code	CSB-PA007770ESR2HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P18074
Immunogen	Recombinant Human TFIIH basal transcription factor complex helicase XPD subunit protein (1-300AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Relevance	ATP-dependent 5'→3' DNA helicase, component of the core-TFIIH basal transcription factor. Involved in nucleotide excision repair (NER) of DNA by opening DNA around the damage, and in RNA transcription by RNA polymerase II by anchoring the CDK-activating kinase (CAK) complex, composed of CDK7, cyclin H and MAT1, to the core-TFIIH complex. Involved in the regulation of vitamin-D receptor activity. As part of the mitotic spindle-associated MMXD complex it plays a role in chromosome segregation. Might have a role in aging process and could play a causative role in the generation of skin cancers.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	General transcription and DNA repair factor IIH helicase subunit XPD (TFIIH subunit XPD) (EC 3.6.4.12) (Basic transcription factor 2 80 kDa subunit) (BTF2 p80) (CXPDP) (DNA excision repair protein ERCC-2) (DNA repair protein complementing XP-D cells) (TFIIH basal transcription factor complex 80 kDa subunit) (TFIIH 80 kDa subunit) (TFIIH p80) (Xeroderma pigmentosum group D-complementing protein), ERCC2, XPD XPDC
Species	Human
Research Area	Epigenetics and Nuclear Signaling
Target Names	ERCC2
Image	



Western blot

All lanes: TFIIH basal transcription factor complex helicase XPD subunit antibody at 2 μ g/ml

Lane 1: 293T whole cell lysate


Lane 2: K562 whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 87, 47 kDa

Observed band size: 87 kDa



Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA007770ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human lung cancer using CSB-PA007770ESR2HU at dilution of 1:100