






## **GRIA3** Antibody

Product CodeCSB-PA009900LA01HUStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P42263ImmunogenRecombinant Human Glutamate receptor 3 protein (151-250AA)	
Uniprot No. P42263	
Immunogen Recombinant Human Glutamate receptor 3 protein (151-250AA)	
Raised In Rabbit	
Species Reactivity Human	
<b>Tested Applications</b> ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:2	00
Receptor for glutamate that functions as ligand-gated ion channel in nervous system and plays an important role in excitatory synaptic trate. L-glutamate acts as an excitatory neurotransmitter at many synapses central nervous system. Binding of the excitatory neurotransmitter L-induces a conformation change, leading to the opening of the cation and thereby converts the chemical signal to an electrical impulse. The then desensitizes rapidly and enters a transient inactive state, charat the presence of bound agonist. In the presence of CACNG4 or CACNG8, shows resensitization which is characterized by a delayed	ensmission. s in the glutamate channel, he receptor cterized by NG7 or
accumulation of current flux upon continued application of glutamate	
• •	
accumulation of current flux upon continued application of glutamate	
accumulation of current flux upon continued application of glutamate  Form  Liquid	
accumulation of current flux upon continued application of glutamate  Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300	
accumulation of current flux upon continued application of glutamate  Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4	
Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method Purification Method	
Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG	3) (GluR-C)
Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Glutamate receptor 3 (GluR-3) (AMPA-selective glutamate receptor (GluR-K3) (Glutamate receptor ionotropic, AMPA 3) (GluA3), GRIA3	3) (GluR-C)
accumulation of current flux upon continued application of glutamate  Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Glutamate receptor 3 (GluR-3) (AMPA-selective glutamate receptor (GluR-K3) (Glutamate receptor ionotropic, AMPA 3) (GluA3), GRIA3 GLURC	3) (GluR-C)
Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Glutamate receptor 3 (GluR-3) (AMPA-selective glutamate receptor (GluR-K3) (Glutamate receptor ionotropic, AMPA 3) (GluA3), GRIA3 GLURC  Species Human	3) (GluR-C)




## **CUSABIO TECHNOLOGY LLC**












Immunohistochemistry of paraffin-embedded human gastric cancer using CSB-PA009900LA01HU at dilution of 1:100



Immunofluorescent analysis of PC-3 cells using CSB-PA009900LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)