


GTF2F2 Antibody

Product Code	CSB-PA010005LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P13984
Immunogen	Recombinant Human General transcription factor IIF subunit 2 protein (2-249AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB; Recommended dilution: WB:1:1000-1:5000
Relevance	TFIIF is a general transcription initiation factor that binds to RNA polymerase II and helps to recruit it to the initiation complex in collaboration with TFIIB. It promotes transcription elongation. This subunit shows ATP-dependent DNA-helicase activity.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	General transcription factor IIF subunit 2 (EC 3.6.4.12) (ATP-dependent helicase GTF2F2) (General transcription factor IIF 30 kDa subunit) (Transcription initiation factor IIF subunit beta) (TFIIF-beta) (Transcription initiation factor RAP30), GTF2F2, RAP30
Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	GTF2F2

Image



Western Blot

All lanes: GTF2F2 antibody at 16μg/ml

Lane 1: PC-3 whole cell lysate

Lane 2: Hela whole cell lysate

Lane 3: K562 whole cell lysate

Lane 4: 293T whole cell lysate

Lane 5: HepG2 whole cell lysate

Lane 6: HL60 whole cell lysate

Lane 7: A549 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 28 kDa

Observed band size: 28 kDa