



HDAC3 Antibody, FITC conjugated

Product Code	CSB-PA010239LC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O15379
Immunogen	Recombinant Human Histone deacetylase 3 protein (1-428AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation.
Form	Liquid
Conjugate	FITC
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Histone deacetylase 3 (HD3) (EC 3.5.1.98) (RPD3-2) (SMAP45), HDAC3
Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	HDAC3