





HDAC8 Antibody, Biotin conjugated

Product Code	CSB-PA010244LD01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9BY41
Immunogen	Recombinant Human Histone deacetylase 8 protein (1-377AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from
	chromatin. May play a role in smooth muscle cell contractility.
Form	Liquid
Form Conjugate	
	Liquid
Conjugate	Liquid Biotin Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Histone deacetylase 8 (HD8) (EC 3.5.1.98), HDAC8, HDACL1