





## **PVR** Antibody

<b>Product Code</b>	CSB-PA019093ESR1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P15151
Immunogen	Recombinant Human Poliovirus receptor protein (220-345AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Relevance	Mediates NK cell adhesion and triggers NK cell effector functions. Binds two different NK cell receptors: CD96 and CD226. These interactions accumulates at the cell-cell contact site, leading to the formation of a mature immunological synapse between NK cell and target cell. This may trigger adhesion and secretion of lytic granules and IFN-gamma and activate cytoxicity of activated NK cells. May also promote NK cell-target cell modular exchange, and PVR transfer to the NK cell. This transfer is more important in some tumor cells expressing a lot of PVR, and may trigger fratricide NK cell activation, providing tumors with a mechanism of immunoevasion. Plays a role in mediating tumor cell invasion and migration. Serves as a receptor for poliovirus attachment to target cells. May play a role in axonal transport of poliovirus, by targeting virion-PVR-containing endocytic vesicles to the microtubular network through interaction with DYNLT1. This interaction would drive the virus-containing vesicle to the axonal retrograde transport.2 Publications
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Poliovirus receptor (Nectin-like protein 5) (NECL-5) (CD antigen CD155), PVR, PVS
Species	Human
Research Area	Microbiology
Target Names	PVR
Image	




## **CUSABIO TECHNOLOGY LLC**









Western blot

All lanes: Poliovirus receptor antibody at 3µg/ml


Lane 1: Mouse gonad tissue Lane 2: HepG2 whole cell lysate Lane 3: K562 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 46, 41, 40, 43 kDa

Observed band size: 46 kDa



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA019093ESR1HU at dilution of 1:100