


Image





SOCS2 Antibody

Product Code	CSB-PA022389ESR2HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O14508
Immunogen	Recombinant Human Suppressor of cytokine signaling 2 protein (1-198AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200
Relevance	SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. SOCS2 appears to be a negative regulator in the growth hormone/IGF1 signaling pathway. Probable substrate recognition component of a SCF-like ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins.
Form	Liquid
Form Conjugate	Liquid Non-conjugated
	•
Conjugate	Non-conjugated
Conjugate Storage Buffer	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate Storage Buffer Purification Method	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified
Conjugate Storage Buffer Purification Method Isotype	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal Suppressor of cytokine signaling 2 (SOCS-2) (Cytokine-inducible SH2 protein 2)
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal Suppressor of cytokine signaling 2 (SOCS-2) (Cytokine-inducible SH2 protein 2) (CIS-2) (STAT-induced STAT inhibitor 2) (SSI-2), SOCS2, CIS2 SSI2 STATI2
Conjugate Storage Buffer Purification Method Isotype Clonality Alias Species	Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal Suppressor of cytokine signaling 2 (SOCS-2) (Cytokine-inducible SH2 protein 2) (CIS-2) (STAT-induced STAT inhibitor 2) (SSI-2), SOCS2, CIS2 SSI2 STATI2 Human



Western blot


All lanes: Suppressor of cytokine signaling 2 antibody at $2\mu g/ml + HepG2$ whole cell lysate Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 22 kDa Observed band size: 22 kDa




CUSABIO TECHNOLOGY LLC












Immunohistochemistry of paraffin-embedded human lung tissue using CSB-PA022389ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA022389ESR2HU at dilution of 1:100