





## UBE2C Antibody, HRP conjugated

Product Code         CSB-PA025440LB01HU           Storage         Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.           Uniprof No.         O00762           Immunogen         Recombinant Human Ubiquitin-conjugating enzyme E2 C protein (2-179AA)           Raised In         Rabbit           Species Reactivity         Human           Tested Applications         ELISA           Relevance         Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-11\'- and 'Lys-48\'- linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'Lys-11\'- linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.           Form         Liquid           Conjugate         HRP           Storage Buffer         Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4           Purification Method         >95%, Protein G purified           Isotype         IgG           Clonality         Polyclonal           Alias         Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (Ubiquitin-conjugating enzyme C) (Ubiquitin-co		
Uniprot No. O00762  Immunogen Recombinant Human Ubiquitin-conjugating enzyme E2 C protein (2-179AA)  Raised In Rabbit  Species Reactivity Human  Tested Applications ELISA  Relevance Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'L'ys-11\'- and 'L'ys-48\'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'L'ys-11\'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.  Form Liquid  Conjugate HRP  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (Ubiquitin-protein ligase C), UBE2C, UBCH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	<b>Product Code</b>	CSB-PA025440LB01HU
Immunogen         Recombinant Human Ubiquitin-conjugating enzyme E2 C protein (2-179AA)           Raised In         Rabbit           Species Reactivity         Human           Tested Applications         ELISA           Relevance         Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-11\'- and 'Lys-48\'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'Lys-11\'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.           Form         Liquid           Conjugate         HRP           Storage Buffer         Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4           Purification Method         >95%, Protein G purified           Isotype         IgG           Clonality         Polyclonal           Alias         Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) ((EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10           Species         Homo sapiens (Human)           Research Area         Epigenetics and Nuclear Signaling	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit  Species Reactivity Human  Tested Applications ELISA  Relevance Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'LLys-11\'- and 'LLys-48\'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'Llys-11\'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.  Form Liquid  Conjugate HRP  Storage Buffer Preservative: 0.03% Proclin 300	Uniprot No.	O00762
Species Reactivity         Human           Tested Applications         ELISA           Relevance         Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-11'- and 'Lys-48\'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating 'Lys-11\'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.           Form         Liquid           Conjugate         HRP           Storage Buffer         Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4           Purification Method         >95%, Protein G purified           Isotype         IgG           Clonality         Polyclonal           Alias         Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbCH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10           Species         Homo sapiens (Human)           Research Area         Epigenetics and Nuclear Signaling	Immunogen	Recombinant Human Ubiquitin-conjugating enzyme E2 C protein (2-179AA)
Tested Applications  Relevance  Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes \(^1\text{Lys-11\'}\)- and \(^1\text{Lys-48\'}\)-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating \(^1\text{Lys-11\'}\)-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.  Form  Liquid  Conjugate  HRP  Storage Buffer  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method  >95%, Protein G purified  Isotype  IgG  Clonality  Polyclonal  Alias  Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species  Homo sapiens (Human)  Research Area  Epigenetics and Nuclear Signaling	Raised In	Rabbit
Relevance Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes \Lys-11\\- and \\Lys-48\\-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating \\Lys-11\\-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.  Form Liquid Conjugate HRP Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG Clonality Polyclonal Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling	Species Reactivity	Human
other proteins. In vitro catalyzes \Lys-11\\- and \Lys-48\\-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating \Lys-11\\-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by the proteasome and promoting mitotic exit.  Form Liquid  Conjugate HRP  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Tested Applications	ELISA
ConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasUbiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear Signaling	Relevance	other proteins. In vitro catalyzes \'Lys-11\'- and \'Lys-48\'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis. Acts by initiating \'Lys-11\'-linked polyubiquitin chains on APC/C substrates, leading to the degradation of APC/C substrates by
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Isotype IgG  Clonality Polyclonal  Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Conjugate	HRP
Clonality Polyclonal  Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Storage Buffer	
Clonality Polyclonal  Alias Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Purification Method	>95%, Protein G purified
Alias  Ubiquitin-conjugating enzyme E2 C (EC 2.3.2.23) ((E3-independent) E2 ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species  Homo sapiens (Human)  Research Area  Epigenetics and Nuclear Signaling	Isotype	IgG
ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C, UBCH10  Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling	Clonality	Polyclonal
Research Area Epigenetics and Nuclear Signaling	Alias	ubiquitin-conjugating enzyme C) (EC 2.3.2.24) (E2 ubiquitin-conjugating enzyme C) (UbcH10) (Ubiquitin carrier protein C) (Ubiquitin-protein ligase C), UBE2C,
	Species	Homo sapiens (Human)
Target Names UBE2C	Research Area	Epigenetics and Nuclear Signaling
	Target Names	UBE2C