


# TNFRSF10D Antibody

<b>Product Code</b>	CSB-PA031451
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9UBN6
<b>Immunogen</b>	Synthesized peptide derived from internal of Human TNFRSF10D.
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Specificity</b>	The antibody detects endogenous levels of total TNFRSF10D protein.
<b>Tested Applications</b>	ELISA, WB; WB: 1:500-1:3000
<b>Relevance</b>	Receptor for the cytotoxic ligand TRAIL. Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis. Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to it cannot but according to it can induce the NF-kappa-B pathway.
	Marsters S.A., Curr. Biol. 7:1003-1006(1997). Degli-Esposti M.A., Immunity 7:813-820(1997). Pan G., FEBS Lett. 424:41-45(1998).
<b>Form</b>	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification Method</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Tumor necrosis factor receptor superfamily member 10D; Decoy receptor 2; DcR2; TNF-related apoptosis-inducing ligand receptor 4; TRAIL receptor 4
<b>Product Type</b>	Polyclonal Antibody
<b>Species</b>	Homo sapiens (Human)
<b>Target Names</b>	TNFRSF10D

**Image**


Western blot analysis of extracts from Jurkat cells, using TNFRSF10D antibody.