


Image






DKK3 Antibody

oduct Code	CSB-PA037884
orage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
niprot No.	Q9UBP4
nmunogen	Fusion protein of Human DKK3
aised In	Rabbit
pecies Reactivity	Human, Mouse
ested Applications	ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:200-1:1000,IHC:1:50-1:200
elevance	This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. The expression of this gene is decreased in a variety of cancer cell lines and it may function as a tumor suppressor gene. Alternative splicing results in multiple transcript variants encoding the same protein.
orm	Liquid
onjugate	Non-conjugated
orage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
urification Method	Antigen affinity purification
otype	IgG
ias	dickkopf WNT signaling pathway inhibitor 3
pecies	Homo sapiens (Human)
arget Names	DKK3
pecies Reactivity ested Applications elevance orm onjugate orage Buffer urification Method otype ias pecies	Human,Mouse ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:200-1:1000,IHC:1:50-1:200 This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryoni development through its interactions with the Wnt signaling pathway. The expression of this gene is decreased in a variety of cancer cell lines and it may function as a tumor suppressor gene. Alternative splicing results in multiple transcript variants encoding the same protein. Liquid Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification IgG dickkopf WNT signaling pathway inhibitor 3 Homo sapiens (Human)



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA037884(DKK3 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: x200)



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA037884(DKK3 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)




CUSABIO TECHNOLOGY LLC









Gel: 8%SDS-PAGE, Lysate: 40 μg, Lane: Human fetal brain tissue, Primary antibody: CSB-PA037884(DKK3 Antibody) at dilution 1/200, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 40 seconds