



🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🧉

INA Antibody

| CSB-PA056378 |
|---|
| Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Q16352 |
| Synthetic peptide of Human INA |
| Rabbit |
| Human,Mouse |
| ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100 |
| Neurofilaments are type IV intermediate filament heteropolymers composed of light, medium, and heavy chains. Neurofilaments comprise the axoskeleton and they functionally maintain the neuronal caliber. They may also play a role in intracellular transport to axons and dendrites. This gene is a member of the intermediate filament family and is involved in the morphogenesis of neurons. |
| Liquid |
| Non-conjugated |
| -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol |
| Antigen affinity purification |
| IgG |
| Homo sapiens (Human) |
| INA |
| |
| |



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA056378(INA Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA056378(INA Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x200)

1



🕜 Tel: +1-301-363-4651 🛛 🖾 Email: cusabio@cusabio.com 🥝 Website: www.cusabio.com 🧉



Gel: 10%SDS-PAGE, Lysate: 40 μg, Lane: Mouse brain tissue, Primary antibody: CSB-PA056378(INA Antibody) at dilution 1/550, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 3 seconds