


Image





HNRNPH2 Antibody

| Product Code | CSB-PA061816 |
|---|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P55795 |
| Immunogen | Synthesized peptide derived from internal of Human hnRNP H. |
| Raised In | Rabbit |
| Species Reactivity | Human,Mouse,Rat |
| Specificity | The antibody detects endogenous levels of total hnRNP H protein. |
| Tested Applications | ELISA,WB;WB:1:500-1:3000 |
| Relevance | This protein is a component of the heterogeneous nuclear ribonucleoprotein (hnRNP) complexes which provide the substrate for the processing events that pre-mRNAs undergo before becoming functional, translatable mRNAs in the cytoplasm. Binds poly(RG). |
| | Vorechovsky I., Genomics 21:517-524(1994). Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). |
| Form | Ross M.T., Nature 434:325-337(2005). |
| Form Purification Method | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, |
| | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity- |
| Purification Method | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Purification Method Clonality | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal |
| Purification Method Clonality Alias | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FTP-3; FTP3; HNRH2; HNRPH2; Heterogeneous nuclear ribonucleoprotein H |
| Purification Method Clonality Alias Product Type | Ross M.T., Nature 434:325-337(2005). Olsen J.V., Cell 127:635-648(2006). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FTP-3; FTP3; HNRH2; HNRPH2; Heterogeneous nuclear ribonucleoprotein H Polyclonal Antibody |



Western blot analysis of extracts from Jurkat cells, 293 cells and K562 cells, using hnRNP H antibody.