

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🤌 Website: www.cusabio.com 🍯

AIFM1 Antibody

Product Code	CSB-PA068462
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O95831
Immunogen	Fusion protein of Human AIFM1
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:1000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100
Relevance	This gene encodes a flavoprotein essential for nuclear disassembly in apoptotic cells, and it is found in the mitochondrial intermembrane space in healthy cells. Induction of apoptosis results in the translocation of this protein to the nucleus where it affects chromosome condensation and fragmentation. In addition, this gene product induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. Mutations in this gene cause combined oxidative phosphorylation deficiency 6, which results in a severe mitochondrial encephalomyopathy. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 10.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	lgG
Alias	apoptosis-inducing factor, mitochondrion-associated, 1
Species	Homo sapiens (Human)
Target Names	AIFM1

Image



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA068462(AIFM1 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x200)

1



🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🥣



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA068462(AIFM1 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: x200)



Gel: 10+12%SDS-PAGE, Lysate: 30 μg, Lane 1-3: NIH/3T3 cells, Hela cells, 293T cells, Primary antibody: CSB-PA068462(AIFM1 Antibody) at dilution 1/600, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds