




CCL8 Antibody


| Product Code | CSB-PA07629A0Rb |
|---------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P80075 |
| Immunogen | Recombinant Human C-C motif chemokine 8 protein (24-99AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC; Recommended dilution: IHC:1:20-1:200 |
| Relevance | Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin. The processed form MCP-2(6-76) does not show monocyte chemotactic activity, but inhibits the chemotactic effect most predominantly of CCL7, and also of CCL2 and CCL5 and CCL8 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | C-C motif chemokine 8 (HC14) (Monocyte chemoattractant protein 2) (Monocyte chemotactic protein 2) (MCP-2) (Small-inducible cytokine A8) [Cleaved into: MCP-2(6-76)], CCL8, MCP2 SCYA10 SCYA8 |
| Species | Human |
| Research Area | Immunology |
| Target Names | CCL8 |
| Image | Immunohistochemistry of paraffin-embedded |



Immunohistochemistry of paraffin-embedded human gastric cancer using CSB-PA07629A0Rb at dilution of 1:100




CUSABIO TECHNOLOGY LLC











Immunohistochemistry of paraffin-embedded human small intestine tissue using CSB-PA07629A0Rb at dilution of 1:100