





MYL2 Antibody, FITC conjugated

| Product Code | CSB-PA10059C0Rb |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P10916 |
| Immunogen | Recombinant Human Myosin regulatory light chain 2, ventricular/cardiac muscle isoform protein (8-164AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA |
| Relevance | Contractile protein that plays a role in heart development and function. Following phosphorylation, plays a role in cross-bridge cycling kinetics and cardiac muscle contraction by increasing myosin lever arm stiffness and promoting myosin head diffusion; as a consequence of the increase in maximum contraction force and calcium sensitivity of contraction force. These events altogether slow down myosin kinetics and prolong duty cycle resulting in accumulated myosins being cooperatively recruited to actin binding sites to sustain thin filament activation as a means to fine-tune myofilament calcium sensitivity to force. During cardiogenesis plays an early role in cardiac contractility by promoting cardiac myofibril assembly. |
| Form | Liquid |
| Conjugate | FITC |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Myosin regulatory light chain 2, ventricular/cardiac muscle isoform (MLC-2) (MLC-2v) (Cardiac myosin light chain 2) (Myosin light chain 2, slow skeletal/ventricular muscle isoform) (MLC-2s/v) (Ventricular myosin light chain 2), MYL2, MLC2 |
| Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Nescardii Area | Signal Transduction |