






MDK Antibody

Product Code Storage Uniprot No. Immunogen Raised In Species Reactivity Tested Applications Relevance	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. P21741 Recombinant Human Midkine protein (21-143AA) Rabbit Human ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated protein kinase (MAPK) and PI3-kinase, and the induction of cell proliferation.
Uniprot No. Immunogen Raised In Species Reactivity Tested Applications	P21741 Recombinant Human Midkine protein (21-143AA) Rabbit Human ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
Immunogen Raised In Species Reactivity Tested Applications	Recombinant Human Midkine protein (21-143AA) Rabbit Human ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
Raised In Species Reactivity Tested Applications	Human ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
Species Reactivity Tested Applications	Human ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200 Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
	Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
Relevance	(PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated
	Involved in neointima formation after arterial injury, possibly by mediating leukocyte recruitment. Also involved in early fetal adrenal gland development
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Midkine (MK) (Amphiregulin-associated protein) (ARAP) (Midgestation and kidney protein) (Neurite outgrowth-promoting factor 2) (Neurite outgrowth-promoting protein), MDK, MK1 NEGF2
Species	Human
Research Area	Neuroscience
Target Names	MDK
Image	Immunofluorescent analysis of HenG2 cells



Immunofluorescent analysis of HepG2 cells using CSB-PA10574A0Rb at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)




CUSABIO TECHNOLOGY LLC









Immunohistochemistry of paraffin-embedded human small intestine tissue using CSB-PA10574A0Rb at dilution of 1:100