


Image





CDK11A/CDK11B Antibody

Product Code	CSB-PA105838
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	B7ZVY7
Immunogen	Synthetic peptide of Human CDK11A/CDK11B
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:50-1:200
Relevance	Cyclin-dependent kinases (CDKs) are a family of protein kinases first discovered for their role in regulating the cell cycle. They are also involved in regulating transcription, mRNA processing, and the differentiation of nerve cells. They are present in all known eukaryotes, and their regulatory function in the cell cycle has been evolutionarily conserved. CDKs are relatively small proteins, with molecular weights ranging from 34 to 40 kDa, and contain little more than the kinase domain. By definition, a CDK binds a regulatory protein called a cyclin. Without cyclin, CDK has little kinase activity; only the cyclin-CDK complex is an active kinase. CDKs phosphorylate their substrates on serines and threonines, so they are serine-threonine kinases.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Target Names	CDK11A/CDK11B



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA105838(CDK11A/CDK11B Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)















The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using CSB-PA105838(CDK11A/CDK11B Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 6%SDS-PAGE, Lysate: 40 μg, Lane 1-4: Hela cells, hepG2 cells, lovo cells, human colon cancer tissue, Primary antibody: CSB-PA105838(CDK11A/CDK11B Antibody) at dilution 1/750, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute