


Image






TACR2 Antibody

CSB-PA125699
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
P21452
Synthetic peptide of Human TACR2
Rabbit
Human
ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100
This gene belongs to a family of genes that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin neuropeptide substance K, also referred to as neurokinin A.
Liquid
Non-conjugated
-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Antigen affinity purification
IgG
Homo sapiens (Human)
TACR2



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA125699(TACR2 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using CSB-PA125699(TACR2 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)




CUSABIO TECHNOLOGY LLC









Gel: 12%SDS-PAGE, Lysate: 30 μ g, Lane 1-2: K562 cells, hela cells, Primary antibody: CSB-PA125699(TACR2 Antibody) at dilution 1/400, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 30 seconds